



**E Journal
of Cardiovascular
Medicine**

| Volume **5** | Issue **4** |
| October-December **2017** |

www.ejcm.org & www.ejcvsmmed.org

E Journal of Cardiovascular Medicine

Correlations between hematological indicators and other known markers in acute coronary syndromes

*Mehmet Ertürk, Fatma Turhan Çağlar, İsmail Bıyık, Nilgün Işıksaçan, Serkan Yazan,
Mehmet Rifat Yıldırım, Ömer Faruk Baycan, Begüm Özalp, Hüseyin Karakurt*

Successful surgical treatment of a ruptured giant atherosclerotic aneurysm of arcus aorta

Bariş Akça, Nevzat Erdil, Olcay Murat Dişli, Mehmet Cengiz Çolak, Bektaş Battaloğlu

Emergent operation of an extensive pulmonary damage after thoracal gunshot assault

Hamit Serdar Başbuğ, Hakan Göçer, Kanat Özışık



E Journal of Cardiovascular Medicine
is a global e-journal targeting articles on:

- **clinical cardiology,**
- **interventional cardiology,**
 - **arrhythmia,**
- **cardiovascular surgery,**
- **vascular & endovascular surgery,**
- **vascular biology**

Editor-in-Chief

Prof. Öztekin Oto

FESC, FACC / President, Heart and Health Foundation of Turkey / Izmir / Turkey

Asistant Editors

Prof. Ali Kutsal

Sami Ulus Children Hospital Department of
Cardiovascular Sugery / Ankara / Turkey

Prof. Erdem Silistreli

Dokuz Eylül University, Department of Cardiovascular
Sugery / Izmir / Turkey

Prof. Bektaş Battaloğlu

İnönü University, Department of Cardiovascular Sugery
Malatya / Turkey

Dr. Onur Saydam

Karaman State Hospital Cardiovascular Surgery
Karaman / Turkey

Dr. Emre Doğan

Trabzon Ahi Evren Cardiovascular Surgery Hospital
Trabzon / Turkey

Dr. Taylan Adademir

Kartal Koşuyolu Resarch Hospital / Istanbul / Turkey

Owner

© TÜSAV Heart and Health Foundation of Turkey

Administration Office

Şair Eşref Bulvarı, 1402 Sk. No. 2/2 Özbaş Apt.
Alsancak - Izmir / Turkey
Tel: + 90 232 464 19 63 / Fax: +90 232 464 24 70
e-mail: info@oztekinoto.com | info@tksv.org

Publishing Coordinator

Hüseyin Kandemir

huseyin@medikalakademi.com.tr

Publisher

Medikal Akademi Yayıncılık ve Prodüksiyon Tic. Ltd. Sti.
Halaskargazi Cad. No: 172, D: 134 - Şişli / İstanbul
Tel: +90 537 309 29 55, Faks: (0212) 233 90 61
www.medikalakademi.com.tr/hizmetlerimiz

International Scientific Advisory Board

Prof. Harald Kaemmerer

German Heart Centre / Munich / Germany

Prof. Marko Turina

University Hospital of Zurich / Zurich / Switzerland

Prof. Frank W. Selke

Chief of Cardiothoracic Surgery at Brown Medical School
Rhode Island / USA

Prof. Joseph E. Bavaria

Hospital of the University of Pennsylvania
Philadelphia / USA

Prof. Fausto Pinto

Director of Lisbon Cardiovascular Institute / Portugal
& President of the European Society of Cardiology

Prof. Lazar Davidovic

Belgrade Medical School Cardiovascular Surgery
Belgrade / Serbia

Prof. Stephan Schueler

Tyne Freeman Hospital, Department for Cardiothoracic
Surgery Newcastle / United Kingdom

Prof. Piotr Kasprzak

University Hospital Regensburg, Director of Vascular
Surgery / Regensburg / Germany

Prof. Jose Luis Pomar

Hospital Clinico de Barcelona, Department of
Cardiovascular Sugery / Barcelona / Spain

www.ejcm.org
www.ejcvsmmed.org

Prof. Mohamed Moustafa Abdelaal

*Kafrelsheikh University Hospital, Cardiothoracic surgery and
General Director / Kafr El Sheikh / Egypt*

Assoc. Prof. Barış Akça

*Inonu University School of Medicine, Department of
Cardiovascular Surgery / Malatya / Turkey*

Dr. Rezan Aksoy

*Siyami Ersek Training and Research Hospital,
Cardiovascular Surgery / Istanbul / Turkey*

Dr. Şafak Alpat

*Birmingham Children's Hospital Pediatric Cardiovascular
Surgery / Birmingham / UK*

Dr. Mustafa Aldemir

*Kocatepe University, Department of Cardiovascular
Surgery / Afyon / Turkey*

Dr. Elena Zapata-Arriaza

*Hospital Universitario Virgen del Rocío, Instituto de
biomedicina de Sevilla, Vascular Medicine / Sevilla / Spain*

Dr. Mehmet Atay

*Bakırköy Sadi Konuk Research Hospital, Cardiovascular
Surgery / Istanbul / Turkey*

Assoc. Prof. Hakan Aydın

*Sami Ulus in Ankara Training and Research Hospital,
Cardiovascular Surgery / Ankara / Turkey*

Assoc. Prof. Ahmet Çağrı Aykan

*Ahi Evren University of Health Sciences, Thoracic and
Cardiovascular Surgery / Trabzon / Turkey*

Assoc. Prof. Vedat Bakuy

*Bakırköy Sadi Konuk Training and Research Hospital,
Cardiovascular Surgery / Istanbul / Turkey*

Dr. Stefano Bartoli

*ASL Roma2, Cardiovascular Surgery
Rome / Italy*

Assoc. Prof. Elif Börekçi

*Bozok University Research and Application Hospital,
Internal Medicine / Yozgat / Turkey*

Dr. Tomasa Centella

*Hospital Ramón y Cajal, Cardiovascular Surgery
Madrid / Spain*

Assoc. Prof. Ahmet Çalışkan

*Dicle University School of Medicine, Cardiovascular Surgery
Diyarbakır / Turkey*

Dr. Gökhan Çavuşoğlu

*Ahi Evren University of Health Sciences, Thoracic and
Cardiovascular Surgery, Radiology / Trabzon / Turkey*

Dr. Deniz Çevirme

*Kartal Koşuyolu Research and Education
Hospital, Cardiovascular Surgery / Istanbul / Turkey*

Prof. Ferit Çiçekçioğlu

*Bozok University, Training and Research Hospital,
Cardiovascular Sugery / Yozgat / Turkey*

Assoc. Prof. Ertan Demirdaş

*Bozok University Research and Application Hospital and
Cardiovascular Surgery / Yozgat / Turkey*

Assoc. Prof. Yüksel Dereli

*Necmettin Erbakan University, Meram Midical Faculty
Hospital, Cardiovascular Surgery / Konya / Turkey*

Assist.Prof. İnci Selin Doğan

*Karadeniz Technical University Faculty of Pharmacy
Pharmacology, Medicinal Chemistry / Trabzon / Turkey*

Dr. Vehbi Doğan

*Sami Ulus Training and Research Hospital, Pediatric
Cardiology / Ankara / Turkey*

Dr. Çağrı Düzyol

*Kocaeli Derince Education and Research Hospital
Cardiovascular Surgery / Kocaeli / Turkey*

Assoc. Prof. Hüseyin Ede

*Bozok University, Medical Faculty, Cardiovascular Surgery
Yozgat / Turkey*

Dr. İlker Ertuğrul

*Sami Ulus Training and Research Hospital, Pediatric
Cardiology / Ankara / Turkey*

Prof. Niyazi Görmüş

*Necmettin Erbakan University, Meram Medical Faculty
Hospital, Cardiovascular Surgery / Konya / Turkey*

Assist. Prof. Adem Güler

*Gulhane Military Medical Academy Department of
Cardiovascular Surgery / Ankara / Turkey*

Assoc. Prof. Mustafa Gülgün

*GATA Department of Pediatrics, Division of Pediatric
Cardiology / Ankara / Turkey*

Prof. Usama Ali M. Hamza

*Mansoura University Faculty of Medicine, Cardiothoracic
Surgery, Cardiovascular Surgery / Mansoura / Egypt /*

Dr. James B Hermiller

*St Vincent's Medical Group, Interventional Cardiology
Indianapolis / USA*

Dr. Akihiko Ikeda

*Tsukuba Medical Center Hospital, Cardiovascular
Surgery / Tsukuba / Japan*

Dr. Richard W Issitt

*Great Ormond Street Hospital, Cardiac Surgery -
Pediatric Cardiology / London / UK*

Dr. Mehmet Kalender

*Derince Training and Research Hospital, Cardiovascular
Surgery / Kocaeli / Turkey*

Dr. Ayşegül Karadeniz

*Ahi Evren University of Health Sciences, Thoracic and
Cardiovascular Surgery, Radiobiology / Trabzon / Turkey*

Assoc. Prof. Osman Kayapınar

*Duzce University, Medical Faculty Department of
Cardiology / Düzce / Turkey*

Assoc. Prof. Alper Kepez

*Marmara University Training and Research Hospital Cardiol-
ogy Clinic / Istanbul / Turkey*

Assoc. Prof. Yasir Khan Khan

*Ch. Pervaiz Elahi Institute of Cardiology, Cardiovascular Sur-
gery / Punjab / Pakistan*

Assoc. Prof. Levent Korkmaz

*Ahi Evren University of Health Sciences, Thoracic and
Cardiovascular Surgery / Trabzon / Turkey*

Assoc. Prof. Ulaş Kumbasar

*Hacettepe University Medical School Cardiovascular
Surgery / Ankara / Turkey*

Dr. Redha Lakehal

*Department of heart surgery, EHS Erriadh / Constantine
Algeria*

Dr. Wei-Chieh Lee

*Kaohsiung Chang Gung Memorial Hospital, Cardiology
Kaohsiung City / Tayvan*

Dr. José Luis Serrano Martínez

*University Hospital of Granada, Department of Internal
Medicine / Granada / Spain*

Assoc. Prof. Ümit Mentеше

*Ahi Evren University of Health Sciences, Thoracic and
Cardiovascular Surgery / Trabzon / Turkey*

Dr. Nooredin Mohammadi

*Iran University of Medical Sciences, Cardiology, Demand
for Health Care, Determinants of Health / Tehran / Iran*

Dr. Vinod Namana

*Maimonides Medical Center, Department of Medical
Research / New York / USA*

Dr. Silvio Nocco

Sirai Hospital, Department of Cardiology / Carbonia / Italy

Assoc. Prof. Zeynep Tuğba Özdemir

*Bozok University School of Medicine, Internal Medicine
Yozgat / Turkey*

Dr. Tanıl Özer

*Kartal Koşuyolu Yüksek İhtisas Research and Education
Hospital / Istanbul / Turkey*

Prof. Murat Özeren

*Mersin University Medical School, Cardiovascular
Surgery / Mersin / Turkey*

Assoc. Prof. Emre Özker

*Başkent University School of Medicine, Department of
Cardiovascular Surgery / Ankara / Turkey*

Dr. Abdullah Özyurt

*Mersin Maternity and Children Diseases Hospital, Pediatric
Cardiology / Mersin / Turkey*

Dr. Recep Oktay Peker

*Hacettepe University, Department of Cardiovascular Surgery
Ankara / Turkey*

Dr. Hikmet Sahratov

*Gülhane Education and Research Hospital, Department of
Cardiovascular Surgery / Ankara / Turkey*

Dr. Gonzalo Luis Alonso Salinas

*Marcelo Sanmartín of Hospital Universitario Ramón y Cajal
Madrid / Spain*

Dr. Stefano Salizzoni

*Città della Salute e della Scienza, Cardiac Surgery,
Cardiac Surgery / Turin / Italy*

Dr. Gökhan Sargın

*Adnan Menderes University Medical School, Internal
Medicine / Aydın / Turkey*

Dr. Mustafa Seren

*Ankara 29 Mayıs State Hospital and Cardiovascular
Surgery / Ankara / Turkey*

Prof. Erdem Silistreli

*Dokuz Eylül University, Department of Cardiovascular
Surgery / İzmir / Turkey*

Assoc. Prof. Ömer Tanyeli

*Necmettin Erbakan University, Meram Medical Faculty
Hospital, Cardiovascular Surgery / Konya / Turkey*

Dr. İlker Tekin

*Antalya Medicalpark Hospital, Cardiovascular Surgery
Antalya / Turkey*

Assist. Prof. Dinçer Uysal

*Isparta Süleyman Demirel University, Department of
Cardiovascular Surgery / Isparta / Turkey*

Dr. Olivier Villemain

*IM3C Necker-Enfants Malades, AP-HP, Université Paris
Descartes, Pediatric Cardiology, Radiology / Paris / France*

Dr. Mustafa Esat Yamaç

*Ahi Evren University of Health Sciences, Thoracic and
Cardiovascular Surgery / Trabzon / Turkey*

Assoc. Prof. Ali Ümit Yener

*Canakkale Onsekiz Mart University Medical Faculty,
Department of Cardiovascular Surgery / Çanakkale / Turkey*

Dr. Dilek Yeşilbursa

*Uludağ University, Medical Faculty, Department of
Cardiology / Bursa / Turkey*

Dr. Mustafa Yılmaz

*Sami Ulus Training and Research Hospital, Pediatric
Cardiology / Ankara / Turkey*

Research Articles

Correlations between hematological indicators and other known markers in acute coronary syndromes | 67

Mehmet Ertürk, Fatma T. Çağlar, İsmail Bıyık, Nilgün Işıksaçan, Serkan Yazan,
Mehmet R. Yıldırım, Ömer F. Baycan, Begüm Ozalp, Hüseyin Karakurt

Case Report

Successful surgical treatment of a ruptured giant atherosclerotic aneurysm of arcus aorta | 75

Barış Akça, Nevzat Erdil, Olcay Murat Dişli, Mehmet Cengiz Çolak, Bektaş Battaloğlu

Combined system usage for large neck anatomy in a high risk abdominal aortic aneurysm patient | 79

Funda Yıldırım, Dilşad Amanvermez Şenarslan, Adnan Taner Kurdal,
Tülün Öztürk, Barış Bayram, Mustafa Cerrahoğlu, İhsan İşkesen, Ömer Tetik

Emergent operation of an extensive pulmonary damage after thoracal gunshot assault | 83

Hamit Serdar Başbuğ, Hakan Göçer, Kanat Özışık

Which should be treated first; coronary artery disease or celiac artery compression syndrome? | 87

İsmail Bıyık, Erol Bahtiyar, Nilgün Işıksaçan

Short Communication

It's not fair to compare transcatheter aortic valve implantation with surgical replacement | 91

Taylan Adademir, Davut Çekmecelioğlu, Deniz Çevirme, Mehmet Kalender,
Kamil Boyacioğlu, Fuat Büyükbayrak, Hızır Mete Alp

Correlations between hematological indicators and other known markers in acute coronary syndromes

Mehmet Ertürk¹, Fatma N. Turhan Caglar², İsmail Bıyık³, Nilgun Işıksacan⁴, Serkan Yazan¹, Mehmet Rifat Yıldırım¹, Omer Faruk Baycan¹, Begum Ozalp¹, Huseyin Karakurt¹

¹⁾ Department of Cardiology, Mehmet Akif Ersoy Thoracic and Cardiovascular Surgery Training and Research Hospital, Istanbul, Turkey

²⁾ Department of Cardiology, Bakirkoy Dr. Sadi Konuk Education and Research Hospital, Istanbul, Turkey

³⁾ Department of Cardiology, Usak University, School of Medicine, Education and Research Hospital, Usak, Turkey

⁴⁾ Department of Biochemistry, Bakirkoy Dr. Sadi Konuk Education and Research Hospital, Istanbul, Turkey

Abstract

Introduction: Recently, many hematologic markers have identified as prognostic and diagnostic indicators in different acute coronary syndrome (ACS) patients. In particular, neutrophil / lymphocyte ratio (NLR) and platelet / lymphocyte ratio (PLR) are recognized as markers in the diagnosis and prognosis of ACS. In this study, our aim was to investigate the correlation between the diagnostic yield of PLR and NLR values and other markers such as troponin in all ACS patients.

Material and methods: 319 patients admitted to our hospital with ACS and 283 control patients were included in the study. Leukocyte, neutrophil, platelet, lymphocyte counts, PLR, NLR and high sensitive troponin I (HsTnI) measurements were taken.

Results: Leukocyte, neutrophil and platelet counts were significantly higher in the ACS group than the controls ($p < 0.001$). Lymphocyte count was significantly lower in the ACS group than the controls ($p < 0.001$). NLR and PLR were found to be significantly higher in the ACS group than the controls (4.0 ± 3.3 vs 2.1 ± 1.4 and 126.3 ± 68.9 vs 106.9 ± 49.4 , $p < 0.001$, respectively). NLR was showed significant correlation with HsTnI, PLR, angina time, presence of ST elevation and T wave negativity ($p < 0.05$), and PLR revealed significant correlation with NLR, HsTnI, ST elevation and T wave negativity ($p < 0.05$).

Conclusion: Hematologic markers were found to be significantly higher in the ACS group than the controls. The difference also continued in subgroup analyzes. NLR and PLR correlated with many other ACS indicators.

Key words: Platelet lymphocyte ratio, neutrophil lymphocyte ratio, acute coronary syndrome, troponin.

Ertürk M., Turhan Caglar FN., Bıyık İ., et al. Correlations between hematological indicators and other known markers in acute coronary syndromes. EJCM 2017; 05 (4): 67-74. Doi: 10.15511/ejcm.17.00467.

Introduction

Coronary artery disease (CAD) is a major cause of morbidity and mortality worldwide.⁽¹⁾ One of the most important causes of CAD is atherosclerosis and atherosclerosis is considered to be an inflammatory disease.⁽²⁾ Previous studies have shown serious evidence that atherosclerosis is systemic rather than focal disease.⁽²⁾ Recently, many studies have been published, which demonstrate that hematological markers measured in peripheral blood count such as platelet, neutrophil, lymphocyte are predictors in the diagnosis and the determination of the prognosis of acute coronary syndromes (ACS).^(3, 4) Increased platelet activation has been reported to play a role in the onset and the progression of atherosclerosis.⁽⁵⁾ It has been reported that increased platelet count is associated with platelet activation and may be associated with thrombotic tendency and major adverse cardiac events.^(3, 5)

On the other hand, there are reports that low lymphocyte counts are associated with lymphocyte apoptosis in prolonged inflammation, decreased immune-reactivity, and may also cause major adverse cardiac events.⁽⁵⁻⁷⁾ As a result, proliferation and relative thrombocytosis are seen in the megakaryocytic series with inflammation.⁽⁷⁾ Thus, in some studies, high platelet / lymphocyte ratio (PLR) and neutrophil / lymphocyte ratio (NLR) are predicted to better reflect inflammation and pro-thrombotic states and may be predictors of prognosis in acute ACS patients.^(7, 8) In our study, we aimed to investigate the diagnostic efficiency of PLR and NLR values and their correlation with other predictors such as high sensitivity troponin I in ACS patients admitted to our hospital.

Patients and Method

This single-center comparative study was conducted in a tertiary heart center. The study was approved by the local Ethics committee and informed consent of the patients and controls were obtained. 319 patients admitted to our hospital with the diagnosis ACS between 2015 and 2017 were included in the study. In the same time period, 283 healthy individuals who applied to our outpatient clinics were included in the study as a control group. Subsequently, the ACS group was divided into

subgroups as unstable angina pectoris (USAP), non-ST elevation myocardial infarction (NSTEMI), and ST elevation myocardial infarction (STEMI). Patients with severe valve disease, decompensated heart failure, malignancy, hematological disease, systemic inflammatory disease, active infection, autoimmune disease, severe renal or hepatic insufficiency and those using steroids were excluded from the study.

The demographic and biochemical parameters of the patients and controls were obtained from the files. NLR and PLR were calculated from the neutrophil, platelet and lymphocyte counts obtained from the hemogram. PLR and NLR were calculated by dividing platelet count by lymphocyte count, respectively, by dividing the number of neutrophils by the number of lymphocytes. For the measurement of high sensitive troponin I (HsTnI) EDTA whole blood sample was used, HsTnI was measured by the method of the AQT90 FLEX cardiac Troponin I immunoassay (Radiometer Medical ApS, Copenhagen, Denmark), and the method is a one-step sandwich immunofluorometric assay based on the use of three monoclonal antibodies, two for the capture and one for the detection.⁽⁹⁾

Complete blood count (CBC) was measured from EDTA whole blood samples via BC 6800 auto analyzer (Mindray Medical International Limited, Shenzhen, China). Other biochemical measurements such as glucose, HDL, LDL, total cholesterol and triglycerides were made from serum samples via Cobas systems (Roche Diagnostic Basel, Switzerland) by using commercial kits (Roche Diagnostic Basel, Switzerland).⁽⁹⁾ Two levels of internal quality controls were made for all devices.

Statistical analysis

Statistical analysis was performed using SPSS version 15.0 (SPSS Inc., Chicago, Illinois, USA). Continuous variables were described by mean and standard deviation. Categorical variables were described in percent. Mann-Whitney U-test and student t-test were used for continuous variables, while chi-square test was used for categorical variables. Kruskal-Wallis test and Mann-Whitney U-test were used for subgroup analysis. Spearman test was used for correlation analysis. The

diagnostic accuracy of high sensitive troponin I, NLR and PLR for ACS was calculated by receiver operating characteristic (ROC) analysis.

Results

319 ACS patients (219, 68.7%, male) and 283 controls (181, 64%, male) were included in the study. There were 114 (35.7%) patients with USAP, 101 (31.7%) patients with NSTEMI and 104 (32.6%) patients with STEMI in the ACS subgroups. The demographic, clinical, electrocardiographic and biochemical parameters of the patients are shown in **Table 1**. Neutrophil and platelet counts were significantly higher in the ACS groups than in the control group ($p < 0.001$ respectively). Lymphocyte count was lower in the ACS group than in the control group ($p < 0.001$). NLR and PLR were significantly higher in the ACS group than in the control group ($p < 0.001$).

In the ACS subgroup analysis, NLR and PLR values were significantly higher in STEMI patients than in USAP and NSTEMI patients ($p < 0.001$) (**Table 2**). Correlation analyzes revealed that NLR correlated with HsTnI, PLR, ST elevation, ST depression, T wave negativity and angina duration (**Table 3**). PLR was correlated with NLR, HsTnI, ST elevation and T wave negativity (**Table 3**). Receiver operating characteristic (ROC) analysis results are given in **Table 4**. When the cut-off value for NLR was taken as 2.5, the sensitivity of NLR for the diagnosis of ACS was found 63.6%, specificity 80.2%, negative predictive value 66.1% and positive predictive value 78.6% (95% confidence in-

terval 0.66-0.75). When the cut off value for PLR was taken as 101.2, the sensitivity of PLR for the diagnosis of ACS was determined as 54.9%, specificity, 54.4%, negative predictive value 51.7% and positive predictive value 57.6% (95% confidence interval 0.54-0.63).

Discussion

In recent years, NLR and PLR have become important markers easily measured from peripheral blood and have proven diagnostic and prognostic information in patients with ACS.⁽¹⁰⁾ In our study, NLR and PLR values were found to be higher in all ACS patients than the control group in accordance with the literature. In subgroup analyzes, NLR and PLR values were significantly higher in STEMI patients than NSTEMI and USAP patients. NLR and PLR were correlated with each other and with other parameters which have proven prognostic information in ACS patients such as high sensitive troponin I (**Table 3**). Sezer et al. showed that the number of neutrophils and the mean platelet volume in patients with acute myocardial infarction were associated with reperfusion injury after infarct-related artery occlusion opened.⁽¹⁰⁾ In many inflammatory conditions neutrophils are stimulated to release many cytokines and cytotoxic or proteolytic enzymes.⁽⁴⁾

These enzymes have adverse effects on the ischemic heart by damaging endothelial cells, stimulating the coagulation system, blocking the microvascular circulation, and increasing infarct expansion.⁽⁴⁾ Lymphocytes, as parts of the adaptive immunity system, fight against inflammation and suppress inflammation.⁽²⁾ Thus, low

Table 2. Subgroups comparisons of hematologic indicators and cardiac markers

	USAP (n=114, 35.7%)	NSTEMI (n=101, 31.7%)	STEMI (n=104, 32.6%)	
	Mean ± Standard Deviation	Mean ± Standard Deviation	Mean ± Standard Deviation	P value
HsTnI (µg/L)	0.0108 ± 0.0084	1.2659 ± 3.6638	5.5973 ± 7.8381	<0.001
NLR	2.61 ± 1.59	3.30 ± 2.36	6.13 ± 4.34	<0.001
PLR	99.27 ± 48.54	125.90 ± 65.59	156.37 ± 78.56	<0.001

HsTnI: High sensitivity troponin I, **NLR:** neutrophil / Lymphocyte ratio, **PLR:** platelet / lymphocyte ratio

Table 1. Demographic characteristics of patients and controls

	Patients (n=319)	Controls (n=283)	P value
Age (years)	56.6 ± 11.4	47.3 ± 13.6	<0.001
Male, n (%)	219 (68.7)	181 (64)	0.223
BMI, kg/m ²	27.8 ± 4.2	27.5±4.3	0.697
History			
Known CAD, n (%)	109 (34.2)	54 (19.1)	<0.001
Previous PCI, n (%)	80 (25.1)	44 (15.5)	0.004
Previous AMI, n (%)	59 (18.5)	32 (11.3)	0.014
Previous CABG, n (%)	51 (16)	13 (4.6)	<0.001
Previous stroke / TIA, n (%)	17 (5.3)	7 (2.5)	0.074
Cardiovascular risk factors, n (%)			
Hypertension, n (%)	145 (45.5)	102 (36)	0.019
Diabetes mellitus, n (%)	57 (17.9)	39 (13.8)	0.172
Family history of CAD, n (%)	80 (25.1)	30 (10.6)	<0.001
Smoking, n (%)	156 (48.9)	127 (44.9)	0.323
Electrocardiographic findings in admission			
Normal, n (%)	87 (27.3)	247 (87.3)	<0.001
ST segment elevation, n (%)	100 (31.3)	0 (0)	
ST segment depression, n (%)	61 (19.1)	6 (2.1)	
T-wave inversion, n (%)	42 (13.2)	24 (8.5)	
Bundle branch block, n (%)	29 (9.1)	6 (2.1)	
Duration of angina, hours	7.9 ± 1.7	8.0 ± 1.8	0.886
Biochemical parameters			
Total cholesterol, mg/dL	202.2 ± 44.3	188.6 ± 42.7	<0.001
LDL cholesterol, mg/dL	140.0 ± 41.6	123.9 ± 37.8	<0.001
HDL cholesterol, mg/dL	44.2 ± 10.5	44.8 ± 14.1	0.321
Triglycerides, mg/dL	167.0 ± 103.1	188.8 ± 125.0	0.046
Glucose, mg/dL	137.4 ± 63.1	115.4 ± 47.1	<0.001
Creatinine, mg/dL	0.96 ± 0.5	0.88 ± 0.47	0.003
High sensitive Troponin I µg/L	2.2295 ± 5.4671	0.0102 ± 0.0054	<0.001
Hemogram parameters			
Hematocrit, %	41.4 ± 5.9	40.6 ± 5.1	0.422
WBC	10.4 ± 3.5	8.5 ± 2.3	<0.001
Neutrophil count 10 ³ / µL	7.2 ± 3.3	5.0 ± 1.9	<0.001
Lymphocyte count 10 ³ / µL	2.3 ± 1.1	2.7 ± 0.9	<0.001
Platelet count 10 ³ / µL	244.5 ± 60.0	260.6 ± 64.8	<0.001
NLR	4.0 ± 3.3	2.1 ± 1.4	<0.001
PLR	126.3 ± 68.9	106.9 ± 49.4	<0.001

BMI: body mass index, **CAD:** coronary artery disease, **AMI:** acute myocardial infarction, **CABG:** coronary artery bypass grafting, **TIA:** transient ischemic attack, **LDL:** low-density lipoprotein cholesterol, **HDL:** high-density lipoprotein cholesterol, **Htc:** hematocrit, **WBC:** white blood cell count, **NLR:** neutrophil / lymphocyte ratio, **PLR:** platelet / lymphocyte ratio.

lymphocyte counts have been associated with atherosclerosis progression and poor clinical outcomes.⁽²⁾ Zouridakis et al found that low lymphocyte counts in unstable angina pectoris patients are strongly associated with future cardiac events.⁽¹¹⁾

Although the pathophysiological mechanism of this condition is not fully understood, lymphocyte count is thought to reflect the early response of myocardial ischemia to physiological stress and systemic immunodeficiency.⁽⁴⁾ Conversely, high lymphocyte counts during

Table 3. Correlation analyzes for NLR and PLR in ACS patients

Spearman's rho	Correlation Coefficient Sig. (2-tailed)	NLR	PLR
NLR	r	1.000	0.733**
	p	.	0.000
	n	319	319
HsTnl	r	0.466**	0.475**
	P	0.000	0.000
	n	319	319
PLR	r	0.733**	1.000
	p	0.000	.
	N	319	319
ST depression	r	-0.183**	0.051
	p	0.001	0.361
	n	319	319
Bundle branch block	r	-0.017	-0.045
	p	0.762	0.422
	n	319	319
Duration of angina	r	0.145**	0.085
	p	0.010	0.129
	n	319	319
Location of infarction	r	-0.023	0.068
	p	0.677	0.227
	n	319	319
T inversion	r	-0.285**	-0.221**
	p	0.000	0.000
	n	319	319
ST segment elevation	r	0.466**	0.365**
	p	0.000	0.000
	n	319	319
Type of angina	r	-0.075	-0.023
	p	0.181	0.676
	n	319	319

HsTnl; high sensitivity troponin I, NLR; neutrophil / Lymphocyte ratio, PLR; platelet / lymphocyte ratio, ACS; acute coronary syndrome

ACS reflect a strong immune response and the prognosis of these patients is better.⁽⁴⁾ Instead of examining neutrophils and lymphocytes separately, NLR has been shown to be more prognostic.⁽²⁾ Wang et al. have shown that NLR is predictive of all-cause mortality and cardiovascular events in patients undergoing coronary angiography.⁽¹²⁾ Likewise, Tamhane et al. found that NLR value in patients with percutaneous coronary intervention is related to in-hospital and 6-month mortality.⁽¹³⁾ Ayca et al. have shown that NLR is associated with stent thrombosis.⁽¹⁴⁾

NLR has been studied by different investigators in relation to different scoring systems showing CAD severity and it has been shown to be related to NLR by SYNTAX, GRACE and TIMI scores.⁽¹⁵⁻¹⁷⁾ In our study, similarly, the NLR values were higher in the ACS group than in the control group. Correlation analyzes showed that NLR correlated with high sensitive troponin I, PLR, ST elevation, ST depression, T wave negativity and angina duration. NLR was found to have a sensitivity of 63.6% and a specificity of 80.2% for ACS diagnosis. The role of platelets in ACS is versatile.⁽⁴⁾ The platelet count is associated with the underlying inflammation because inflammatory mediators stimulate megakaryocytic proliferation and result in a prothrombotic state with relative thrombocytosis.^(4, 5)

As platelet levels increase, more platelet rich thrombosis is observed in atherosclerotic plaques, and anti-platelet therapy response is reduced.⁽⁴⁾ It is known that platelets are associated with endothelial activation as well as inflammatory functions.⁽²⁾ Active platelets release

pro-angiogenic mediators and regulate the microvasculature of blood vessels.⁽¹⁸⁾ As essential components of coronary thrombus formation, platelets play an important role in ACS.⁽²⁾ In the CADILLAC study, increased platelet counts were also associated with restenosis and stent thrombosis.⁽¹⁹⁾ PLR is thought to better reflect both inflammation and coagulation pathways.⁽⁴⁾ In their study using optical coherence tomography, Wang et al showed that high PLR is associated with sensitive plaque properties such as thinner fibrous cap, wider lipid load in non-target vessels in ACS patients.⁽²⁰⁾

The authors suggested that the relation between PLR and sensitive plaques is associated with immunologic and inflammatory pathways.⁽²⁰⁾ The relationship between PLR and CAD severity is also shown in other publications.⁽²¹⁾ For example, Kurtul et al. have shown that high PLR is associated with moderate to high SYNTAX score in patients with ACS.⁽²¹⁾ In another study, high platelet counts have been shown to be associated with microvascular plug formation, thrombus formation, and vasoconstriction and no-reflow development.⁽⁸⁾ PLR has been shown to be associated with mortality in STEMI and NSTMI patients.⁽²²⁾ Oylumlu et al noted that high PLR values in ACS patients were associated with in-hospital mortality.⁽⁴⁾

In their meta-analysis, Li et al. showed that increased PLR is associated with in hospital and long-term mortality and cardiovascular events in ACS patients.⁽⁵⁾ Yıldız et al. have also shown that high PLR values are associated with no-reflow development after percutaneous interventions.⁽²³⁾ Prajapati et al. showed a relationship between

Table 4. Receiver operating characteristic (ROC) analyzes of HsTnI, NLR and PLR for diagnosis of ACS

	AUC	SE	P-value	CI%95		Specificity	Sensitivity	NPV	PPV	Cut off value
				Lower limit	Upper limit					
HsTnI	0.800	0.018	0.000	0.764	0.836	99.6	54.2	65.9	99.4	>0.0023
NLR	0.709	0.022	0.000	0.667	0.751	80.2	63.6	66.1	78.6	>2.5
PLR	0.586	0.023	0.000	0.541	0.632	54.4	54.9	51.7	57.6	>101.2

HsTnI; high sensitivity troponin I, *NLR*; neutrophil / Lymphocyte ratio, *PLR*; platelet / lymphocyte ratio, *ACS*; acute coronary syndrome, *NPV*; negative predictive value, *PPV*; positive predictive value

increased PLR and NLR and decreased HDL values in CAD patients.⁽¹⁸⁾ The combined use of NLR and PLR suggests that more prognostic information will be obtained. In this context, Choi et al. found that combined NLR and PLR elevations were associated with long-term cardiovascular events (22). Hematological markers have also been shown in other publications as indicators of inflammation. Similar to other studies, in our study, NLR and PLR in patients with ACS were higher than in the control group. The difference of present study from the other studies is that all the ACS subgroups were included and examined in the same study and that the NLR and PLR were high in all the subgroups.

Conclusion

Inflammatory and thrombotic processes play a crucial role in the development of atherosclerosis, atherosclerotic plaque destabilization and clot formation after plaque rupture. In the diagnosis and prognosis of ACS, there are still needs for reliable, easy, inexpensive and fast detectable markers. PLR and NLR are cheap and easy to use markers in the diagnosis and the prognosis of ACS patients but definite cut off values are needed to use these markers. Large scale and comprehensive studies are needed to reveal absolute cut off values for these markers.

References

1. Bekler A, Gazi E, Yılmaz M, Temiz A, Altun B, Barutcu A, Peker T. Could elevated platelet-lymphocyte ratio predict left ventricular systolic dysfunction in patients with non-ST elevated acute coronary syndrome? *Anatol J Cardiol* 2015; 15: 385-390.
2. Budzianowski J, Pieszko K, Burchardt P, Hickiewicz J. The role of hematological indices in patients with acute coronary syndrome. *Hindawi Disease Markers* 2017; 2017: 3041565. doi: 10.1155/2017/3041565.
3. Thaulow E, Erikssen J, Sandvik L, Stormorken H, Cohn PF. Blood platelet count and function are related to total and cardiovascular death in apparently healthy men. *Circulation* 1991; 84: 613-617.
4. Oylumlu M, Yıldız A, Polat N, Bilik MZ, Akyuz A, Acet H, Soydisc S. Platelet-to-lymphocyte ratio is a predictor of in-hospital mortality patients with acute coronary syndrome. *Anatol J Cardiol* 2015; 15: 277-283.
5. Li H, Zhou Y, Ma Y, Han S, Zhou L. The prognostic value of the platelet-to-lymphocyte ratio in acute coronary syndrome: a systematic review and meta-analysis. *Kardiologia Polska* 2017; 75: 666-673.
6. Ommen SR, Hammill SC, Gibbons RJ. The relative lymphocyte count predicts death in patients receiving implantable cardioverter defibrillators. *Pacing Clin Electrophysiol* 2002; 25: 1424-1428.
7. Edem E, Türker MP, Erkan A, Taş S, Çil A, Vural MG, Akdeniz B. Platelet-to-lymphocyte ratio but not neutrophil-to-lymphocyte ratio predicts high on-treatment platelet reactivity in clopidogrel-treated patients with acute coronary syndrome. *Indian J Pharmacol* 2016; 48: 355-359.
8. Azab B, Shah N, Akerman M, McGinn JT Jr. Value of platelet/lymphocyte ratio as a predictor of all-cause mortality after non-ST elevation myocardial infarction. *J Thromb Thrombolysis* 2012; 34: 326-334.
9. Isiksacan N, Biyik I, Koser M, Ozalp B, Baycan OF, Yazan S, Aktürk IF. Comparison of high sensitive and conventional troponin assays in diagnosis of acute myocardial infarction. *Turk J Biochem* 2017; 42: 77-85.
10. Sezer M, Okcular I, Goren T, Oflaz H, F, Bilge A, Meric M, Umman S. Association of haematological indices with the degree of microvascular injury in patients with acute anterior wall myocardial infarction treated with primary percutaneous coronary intervention. *Heart* 2007; 93: 313-318.
11. Zouridakis EG, Garcia-Moll X, Kaski JC. Usefulness of the blood lymphocyte count in predicting recurrent instability and death in patients with unstable angina pectoris. *Am J Cardiol* 2000; 86: 449-51.
12. Wang X, Zhang G, Jiang X, Lu Z, Xu L. Neutrophil to lymphocyte ratio in relation to risk of all-cause mortality and cardiovascular events among patients undergoing angiography or cardiac revascularization: a meta-analysis of observational studies. *Atherosclerosis* 2014; 234: 206-213.
13. Tamhane UU, Aneja S, Montgomery D, Eagle KA, Gurm HS. Association between admission neutrophil to lymphocyte ratio and outcomes in patients with acute coronary syndrome. *Am J Cardiol*. 2008; 102: 653-7.
14. Ayça B, Akın F, Celik O, Sahin I, Okuyan E, Dinçkal MH. Neutrophil to lymphocyte ratio is related to stent thrombosis and high mortality in patients with acute myocardial infarction. *Angiology* 2015; 66: 545-552.

15. Altun B, Turkon H, Tasolar, Gazi E, Barutcu A, Bekler A, Colkesen Y. The relationship between high-sensitive troponin T, neutrophil lymphocyte ratio and SYNTAX score. *Scand J Clin Lab Invest.* 2014; 74: 108-15.
16. Ergelen M, Uyarel H, Altay S, Kul S, Ayhan E, Isik T, Erdogan E, Turfan M. Predictive value of elevated neutrophil to lymphocyte ratio in patients undergoing primary angioplasty for ST-segment elevation myocardial infarction. *Clin Appl Thromb Hemost.* 2014; 20: 427-32.
17. Sawant AC, Adhikari P, Narra SR, Srivatsa SV, Mills PK, Srivatsa SS. Neutrophil to lymphocyte ratio predicts short- and long-term mortality following revascularization therapy for ST elevation myocardial infarction. *Cardiol J.* 2014; 21: 500-8.
18. Prajapati JH, Sahoo S, Nikam T, Shah KH, Maheriya B, Parmar M. Association of high density lipoprotein with platelet to lymphocyte and neutrophil to lymphocyte ratios in coronary artery disease patients. *J Lipids.* 2014; 2014: 686791.
19. Nikolsky E, Grines CL, Cox DA, Garcia E, Tchong JE, Sadeghi M, Mehran R, Stone GW. Impact of baseline platelet count in patients undergoing primary percutaneous coronary intervention in acute myocardial infarction (from the CADILLAC trial). *Am J Cardiol.* 2007; 99: 1055-61.
20. Wang X, Xie Z, Liu X, Huang X, Lin J, Yu B, Hou J. Association of platelet to lymphocyte ratio with non-culprit atherosclerotic plaque vulnerability in patients with acute coronary syndrome: an optical coherence tomography study. *BMC Cardiovasc Disord.* 2017; 17: 175.
21. Kurtul A, Murat SN, Cetin M, Ornek E. Association of platelet-to-lymphocyte ratio with severity and complexity of coronary artery disease in patients with acute coronary syndromes. *Am J Cardiol.* 2014; 114: 972-8.
22. Choi SW, Choi DH, Ku YH, Ha SI, Park G. Clinical outcome prediction from mean platelet volume in patients undergoing percutaneous coronary intervention in Korean cohort: Implications of more simple and useful test than platelet function testing. *Platelets.* 2014; 25: 322-327.
23. Yıldız A, Yuksel M, Oylumlu M, Polat N, Akyuz A, Acet H, Aydin M, Ulgen MS. The utility of the platelet-lymphocyte ratio for predicting no reflow in patients with ST-segment elevation myocardial infarction. *Clin Appl Thromb Hemost.* 2015; 21: 223-8.

Received: 29/07/2017

Accepted: 24/11/2017

Published: 15/12/2017

Disclosure and conflicts of interest:

Conflicts of interest were not reported.

Corresponding author:

Dr. İsmail Bıyık / (Tel:+905424173209)

Mail: ismailbiyikmd@yahoo.com

Ertürk M., Turhan Caglar FN., Bıyık İ., et al. Correlations between hematological indicators and other known markers in acute coronary syndromes. EJCM 2017; 05 (4): 67-74. Doi: 10.15511/ejcm.17.00467.

Successful surgical treatment of a ruptured giant atherosclerotic aneurysm of arcus aorta

Barış Akça¹, Nevzat Erdil², Olcay Murat Dişli³, Mehmet Cengiz Çolak², Bektaş Battaloğlu²

¹) Department of Cardiovascular Surgery, İnönü University School of Medicine, Ass. Dr., Malatya, Turkey.

²) Department of Cardiovascular Surgery, İnönü University School of Medicine, Prof. Dr., Malatya, Turkey.

³) Department of Cardiovascular Surgery, İnönü University School of Medicine, Malatya, Asst. Assoc. Dr., Turkey.

Abstract

Endovascular aortic repairs in hybrid theatres have been implemented as an alternative option for the treatment of aortic arch aneurysms. The improvements in synthetic graft technologies, cardiopulmonary bypass techniques, blood protection, myocardial protection and protection of the central nervous system have resulted in safer applications of surgical interventions, particularly to the arcus aorta. In this article, we present our successful emergency surgical therapeutic approach in a 74-year-old female patient case with a ruptured giant aortic arch aneurysm.

Key words: Endovascular aortic repairs, aortic arch aneurysms, arcus aorta.

Akça B., Erdil N., Dişli O. M., Çolak M. C., Battaloğlu B. Successful surgical treatment of a ruptured giant atherosclerotic aneurysm of arcus aorta EJCM 2017; 05 (4): 75-78. Doi: 10.15511/ejcm.17.00475.

Introduction

Due to its high mortality and morbidity rates, the surgical treatment for aortic arch aneurysms is challenging,⁽¹⁾ although there is currently a lack of consensus on the optimal treatment modality in the high-risk patients. Recently, endovascular aortic repairs in hybrid theatres have been implemented as an alternative option for the treatment of aortic arch aneurysms.⁽¹⁾ The improvements in synthetic graft technologies, cardiopulmonary bypass techniques, blood protection, myocardial protection and protection of the central nervous system (particularly selective cerebral perfusion and hypothermic circulatory arrest) have resulted in safer applications of surgical interventions, particularly to the arcus aorta.⁽²⁻⁵⁾ In this article, we present our successful emergency surgical therapeutic approach in a female case with a ruptured giant aortic arch aneurysm.

Case Report


A 74-year-old female patient was admitted to our emergency clinic with complaints of blurred consciousness, chest pain and dyspnea. A physical examination of arterial blood pressure, pulse and oxygen saturation revealed values of 80/40 mmHg, 122 beats/min and 85–90%, respectively. On auscultation of the left hemithorax, breath sounds decreased, and a chest X-ray revealed a giant aortic aneurysm and opacity in the left hemithorax that was suggestive of a pleural effusion (**Figure 1A**).

A true saccular aortic arch aneurysm that was measured 8x6 cm² which had ruptured into the left hemithorax was detected in a contrasted thoracoabdominal tomography (**Figure 2A, 2B**).

Written informed consent was obtained from the patient's relatives, and the patient who underwent emergency surgery. The left lung was adhered to the pleura, and the aortic arch was extremely atherosclerotic. A substantial amount of hematoma was present in the left thorax. A cardiopulmonary bypass was performed using the cannulation of the right axillary artery.

The patient was cooled to 25°C and aneurysmal sac opened by applying circulatory arrest under the right selective antegrade cerebral perfusion. An anastomosis of a 30-mm Hemashield graft (Maquet, France) was applied to the descending aorta, and the branches of the arcus were implanted onto the grafts cutting in the form of islets. The graft was then clamped, and a total bypass was applied. The total aortic arch replacement was completed through the graft anastomosis with the ascending aorta during the heating phase. The cardiopulmonary bypass was uneventful. The incisions were closed after bleeding management, and the patient was transferred to the Intensive Care Unit (ICU). She recorded no hemodynamic or neurological problems in her post-operative follow-up, although she was unable to tolerate extubation during the ICU stay (in the postoperative 10th hour), and she was re-intubated due to pulmonary complications.

Figure 1A - 1B



The patient was then transferred to the Anesthesiology and Reanimation Clinic and was extubated by administering a successful ICU program (**Figure 1B**). She experienced no problems following extubation, and she was discharged on the 15th postoperative day.

Discussion

Aortic ruptures, as a complication of aortic aneurysms, leads to massive bleeding with a high rate of mortality. An emergent diagnosis and treatment following a multidisciplinary approach can lead to successful outcomes, despite high mortality rates, as in our case.


With the developments in technological methods and increased surgical experience over the past two decades, successful therapeutic results have been reported in patients with aortic aneurysms.⁽⁶⁾ Applications of minimally invasive endovascular methods that can be performed via small inguinal incisions rather than thoracic aortic interventions through major surgical incisions have been rapidly increasing in recent years.⁽⁷⁾ Hybrid methods and conventional open surgery have been compared in a study that investigated aortic arch interventions, and higher technical success rates with reduced mortality rates have been reported with the application of hybrid methods in a certain high-risk patient group.⁽¹⁾

Previous reports of hybrid arch procedures showed an early mortality rate of 7.4–23.7% with an incidence of stroke seen in 0–13.1%.^(1,8) In a meta-analysis with 463 patients who underwent hybrid arch surgery, the authors reported a 30-day mortality rate of 8.3% with an incidence rate of stroke and paraplegia of 4.4 and 3.9%, respectively.⁽⁹⁾

Endovascular treatments can be administered in cases of thoracic aortic disease that require a surgical approach if open surgery is considered to be a particularly high risk and if the anatomical appearance is appropriate. Our case was subjected to an open surgical intervention due to her presentation with hemodynamic instability and poor clinical status on emergency admission. Furthermore, there was no opportunity to make an endovascular intervention at the time of her admission, and an urgent intervention was needed.

In previous studies, it has been suggested that no definitive evidence exists that supports the superiority of hybrid thoracic endovascular aortic repair (TEVAR) over open arch repair,⁽¹⁰⁾ although most cases of hybrid TEVAR are in high-risk patients who are ineligible for conventional open aortic repair. In this respect, comparing and interpreting the results can be challenging. Further prospective, randomized controlled trials are required to

Figure 2A-2B



compare the outcomes of both surgical strategies.

In conclusion, although the recent outcomes of open arch repair and hybrid TEVAR demonstrate acceptable results, particularly early after the procedure, open arch

repair can bring more reliable outcomes during follow-up. These two surgical strategies, when properly selected based on the individual risk, can improve surgical outcomes in patients with aortic arch aneurysms, especially in ruptured cases.

References

1. Milewski RK, Szeto WY, Pochettino A, Moser GW, Moeller P, Bavaria JE. Have hybrid procedures replaced open aortic arch reconstruction in high-risk patients? A comparative study of elective open arch debranching with endovascular stent graft placement and conventional elective open total and distal aortic arch reconstruction. *J Thorac Cardiovasc Surg* 2010;140(3):590-7.
2. Bachet J, Guilmet D, Goudot B, Dreyfus GD, Delentdecker P, Brodaty D et al. Antegrade cerebral perfusion with cold blood: a 13-year experience. *Ann Thorac Surg* 1999;67:1874-8.
3. Kazui T, Washiyama N, Muhammad BA, Terada H, Yamashita K, Takinami M et al. Total arch replacement using aortic arch branched grafts with the aid of antegrade selective cerebral perfusion. *Ann Thorac Surg* 2000;70:3-9.
4. Kazui T, Yamashita K, Washiyama N, Terada H, Bashar AH, Suzuki K et al. Aortic arch replacement using selective cerebral perfusion. *Ann Thorac Surg* 2007;83:S796-8.
5. Zierer A, Detho F, Dzemali O, Aybek T, Moritz A, Bakhtiary F. Antegrade cerebral perfusion with mild hypothermia for aortic arch replacement: single-center experience in 245 consecutive patients. *Ann Thorac Surg* 2011;91:1868-74.
6. Coselli JS, LeMaire SA. Tips for successful outcomes for descending thoracic and thoracoabdominal aortic aneurysm procedures. *Semin Vasc Surg* 2008;21:13-20.
7. Donas KP, Czerny M, Guber I, Teufelsbauer H, Nanobachvili J. Hybrid open-endovascular repair for thoracoabdominal aortic aneurysms: current status and level of evidence. *Eur J Vasc Endovasc Surg* 2007;34:528-33.
8. Vallejo N, Rodriguez-Lopez JA, Heidari P, Wheatley G, Caparrelli D, Ramaiah V et al. Hybrid repair of thoracic aortic lesions for zone 0 and I in high-risk patients. *JVasc Surg* 2012;55:318-25.
9. Koullias GJ, Wheatley GH III. State-of-the-art of hybrid procedures for the aortic arch: a meta-analysis. *Ann Thorac Surg* 2010;90:689-97.
10. Benedetto U, Melina G, Angeloni E, Codispoti M, Sinatra R. Current results of open total arch replacement versus hybrid thoracic endovascular aortic repair for aortic arch aneurysm: a meta-analysis of comparative studies. *J Thorac Cardiovasc Surg* 2013;145:305-6.

Received: 29/07/2017

Accepted: 18/11/2017

Published: 15/12/2017

Disclosure and conflicts of interest:

Conflicts of interest were not reported.

Corresponding author:

Dr. Barış Akça

Mail: dr.baris.akca@yandex.com.tr

Combined system usage for large neck anatomy in a high risk abdominal aortic aneurysm patient

Funda Yıldırım¹, Dilşad Amanvermez Şenarşlan¹, Adnan Taner Kurdal², Tülün Öztürk³, Barış Bayram¹, Mustafa Cerrahoğlu⁴, İhsan İşkesen⁴, Ömer Tetik⁴

¹ Manisa Celal Bayar University Hafsa Sultan Hospital Cardiovascular Surgery Department, MD., Manisa, Turkey

² Manisa Celal Bayar University Hafsa Sultan Hospital Cardiovascular Surgery Department, Assist.Prof., Manisa, Turkey

³ Manisa Celal Bayar University Hafsa Sultan Hospital Department of Anesthesiology and Reanimation, Prof.Dr., Manisa, Turkey

⁴ Manisa Celal Bayar University Hafsa Sultan Hospital Cardiovascular Surgery Department, Prof.Dr., Manisa, Turkey

Abstract

Endovascular aneurysm repair (EVAR) is the first line approach in abdominal aortic aneurysm (AAA) repair. Open surgical repair of AAA is associated with high perioperative mortality and morbidity. A sufficient infrarenal aortic neck is one of the key points for successful outcome after EVAR. We presented in this case report a 75 year-old, male patient with AAA which maximal diameter is 60 mm. The aneurysm had a large neck that presents a hostile neck anatomy. He was unfit for conventional surgery because of comorbidities as chronic obstructive pulmonary disease, newly diagnosed lung cancer and coronary artery disease. He applied to the emergency service with severe and acutely onset back pain and he was admitted to our cardiovascular surgery clinic. Because of acute onset of the back pain, his serious comorbidities and we planned to use of thoracic and abdominal aortic endovascular stent-graft systems indwelling each other which is known as Funnel technique. Despite progress in endovascular therapy, large infrarenal necks remain a challenge for endovascular interventionists. It seems feasible to use combined thoracic and abdominal stent graft as an alternative treatment modality when the patient is not suitable for standard EVAR and open surgery.

Key words: Endovascular aneurysm repair, abdominal aortic aneurysm, large neck anatomy.

Yıldırım F., Amanvermez Şenarşlan D., Kurdal A. T., Öztürk T., Bayram B., Cerrahoğlu M., İşkesen İ., Tetik Ö. Combined system usage for large neck anatomy in a high risk abdominal aortic aneurysm patient EJCM 2017; 05 (4): 79-82. Doi: 10.15511/ejcm.17.00479.

Introduction

Nowadays, endovascular aneurysm repair (EVAR) is the first line approach in abdominal aortic aneurysm (AAA) repair. Open surgical repair of AAA is associated with high perioperative mortality and morbidity. EVAR is a less invasive method than conventional surgery. A sufficient infrarenal aortic neck is one of the key points for successful outcome after EVAR. According to computed tomography angiography (CT), preoperative sizing and planning should be done precisely.⁽¹⁾

The neck length, angulation, shape, existence of thrombus and calcification should be determined. When the anatomic features of the aneurysm are not favourable for standard EVAR therapies; it is termed “hostile neck anatomy”. Proximal neck characteristics remain a challenge in endovascular treatments. Patients with hostile neck anatomy can be managed in some alternative ways; open surgery, chimney technique, fenestrated endovascular grafts and also Funnel technique described by Zachetta et al.^(2,3)

Fenestrated endovascular abdominal aortic aneurysm repair (FEVAR) and chimney technique are more expensive, off the shelf fenestrated grafts are not available in emergency and in some countries health insur-

ance may not cover the expense of these advanced techniques. In these circumstances funnel technique in which a thoracic stent-graft is placed through the main body of a bifurcated or uni-iliac endograft, must kept in mind.

Case

We presented in this case report a 75 year- old, male patient with AAA which maximal diameter is 60 mm. The aneurysm had a large neck that presents a hostile neck anatomy. He was unfit for conventional surgery because of comorbidities as chronic obstructive pulmonary disease, newly diagnosed lung cancer and coronary artery disease. He applied to the emergency service with severe and acutely onset back pain and he was admitted to our cardiovascular surgery clinic.

In the CT Angiographic examination there was no dissection but he had an AAA with a large neck (38 mm internal diameter at just below the renal artery level) as shown in **Figure 1, 2, 3**. Because of acute onset of the back pain, his serious comorbidities and also because of insurance policies of our country we planned to use of thoracic and abdominal aortic endovascular stent-graft systems indwelling each other which is known as Funnel technique.

Figure 1. AAA with a large neck (38 mm internal diameter at just below the renal artery level), **axial** view of computed tomography angiography.



Figure 2. AAA with a large neck (38 mm internal diameter at just below the renal artery level), **coronal** view of computed tomography angiography.



He was treated combining Valiant Captivia® (Medtronic, Minneapolis, MN) and Endurant® (Medtronic, Minneapolis, MN) endoprosthesis. We prepared the patient for intervention in hybrid operating room with epidural anaesthesia. Both femoral arteries were surgically exposed for arterial access and snared with purse-string sutures to control bleeding from access sites. Firstly, Valiant Captivia Thoracic Stent Graft® 44-44 mm diameter and 100mm length stent graft was deployed just below the lowest renal artery. Then Endurant II® aorto-uni-iliac 36-14 mm diameter and 102 mm length stent-graft was deployed inside of the previous endograft with an overlap of 50 mm.

Later Endurant II® 16-16 mm diameter and 82 mm length extension stent-graft was inserted to the left iliac arterial system. The right common iliac artery was coiled with endoluminal occluder® (OCL22, Medtronic Cardiovascular, Santa Rosa, CA). The control angiography showed a trace of endoleak from the proximal neck. The Reliant balloon catheter® was used for proximal landing zone fixation. The control angiography didn't show any type I endoleak after ballooning of the proximal neck.

At last, uni-iliac EVAR is accomplished by a left to right femorofemoral bypass operation with an 8-mm ringed PTFE graft. The operation was ended without any trouble. The patient was discharged from the hos-

Figure 3. Preoperative 3-dimensional CT angiography reconstruction; hostile neck and hostile iliac artery anatomy.



pital at postoperative third day. The patient has still no problem in his first and third month visits.

Discussion

As many as 20-50% of AAA patients have anatomy that is not convenient for a standard endovascular therapy.⁽⁴⁾ Funnel technique is an alternative way for patients with large hostile neck anatomy who are not suitable for open surgery and the other endovascular methods as FEVAR and chimney grafts. There are limited number of studies related to combined usage of thoracic and abdominal stent-grafts (Funnel Technique).⁽²⁻⁵⁾

Most of the literature consists of case reports. Zachetta et al. first described the method in a case report in 2006.⁽²⁾ Ronsivalle S. et al. described Funnel technique for EVAR: "a way out" for AAA with ectatic proximal necks.⁽⁴⁾ They both reported that the method was secure and effective in sealing to a large aortic neck. They did not observe type I endoleak or migration over a mid-term follow up.

Many manufacturers latest-generation abdominal stent grafts have maximal 35-36 mm diameter neck for trunk and ipsilateral leg endoprosthesis. But when the patient has a larger neck and given the 10-20% oversizing ratio for adequate sealing, the interventionist might need additional stents for endovascular treatment.

Secure fixation and adequate sealing zone at the proximal and distal AAA necks are one of the most important factors in predicting a successful outcome of EVAR.⁽²⁾ Large stent grafts are more at the risk of proximal endoleak and reintervention in the long run.^(6,7)

The authors emphasized the importance of at least 40 mm overlapping to the thoracic stent-graft, leaving more than 40 mm extending beyond the main body of the bifurcated device to ensure expansion to its nominal diameter and adequate sealing.^(2,4) In our case, we also obeyed this protocol.

In a recent case report and literature review Pecoraro et al. compared large neck (24-34 mm) AAA treated by standard stent-grafts and wide neck (≥ 35 mm) AAA treated by Funnel technique. He reports; surprisingly

higher incidence of migrations in large neck group. Authors interpreted as Funnel technique may be a more valuable deployment methods than we thought.⁽⁵⁾

In some clinics, all patients who undergo EVAR are applied sac thrombization with intrasac biomaterials to prevent endoleaks.^(2,4) But in our country, it is not possible because health insurance does not cover its expense. So, we did not apply sac thrombization. We have not yet mid-term and long-term follow-up results. We need further follow-up reports to be sure of its durability.

References

1. De Vries J-PPM. The proximal neck: the remaining barrier to a complete EVAR world. *Semin Vasc Surg* 2012; 25(4): 182–6.
2. Zanchetta M, Faresin F, Pedon L, Riggi M, Colonna S, Lipari R, et al. Funnel technique for first-line endovascular treatment of an abdominal aortic aneurysm with an ectatic proximal neck. *J Endovasc Ther* 2006; 13(6): 775–8.
3. Massara M, Prunella R, Gerardi P, De Caridi G, Serra R, Notarstefano S, et al. A Combination of Thoracic and Abdominal Stent-Grafts to Treat An Abdominal Aortic Aneurysm with Hostile Proximal Neck. *Ann Vasc Surg* 2017; 39: 292.e5-292.e8.
4. Ronsivalle S, Faresin F, Franz F, Rettore C, Zanchetta M, Zonta L. Funnel technique for EVAR: ‘a way out’ for abdominal aortic aneurysms with ectatic proximal necks. *Ann Vasc Surg* 2012; 26(1): 141–8.
5. Pecoraro F, Pakeliani D, Dinoto E, Bajardi G. Endovascular treatment of large and wide aortic neck: case report and literature review. *Gen Thorac Cardiovasc Surg* 2017; 65(4): 219–24.
6. Gargiulo M, Gallitto E, Wattez H, Verzini F, Bianchini Massoni C, Loschi D, et al. Outcomes of endovascular aneurysm repair performed in abdominal aortic aneurysms with large infrarenal necks. *J Vasc Surg* 2017;3:1-8. DOI: 10.1016/j.jvs.2017.01.066. [Epub ahead of print]
7. Kaladji A, Steintmetz E, Gouëffic Y, Bartoli M, Cardon A, Academic Association for Surgical Research (AURC). Long-Term Results of Large Stent Grafts to Treat Abdominal Aortic Aneurysms. *Ann Vasc Surg* 2015; 29(7): 1416–25.

Received: 16/06/2017

Accepted: 04/11/2017

Published: 15/12/2017

Disclosure and conflicts of interest:

The authors declare no conflict of interest.

Corresponding author:

MD. Funda Yıldırım

Mail: fundanizamoglu@yahoo.com

Emergent operation of an extensive pulmonary damage after thoracal gunshot assault

Hamit Serdar Başbuğ¹, Hakan Göçer², Kanat Özişik³

¹) Department of Cardiovascular Surgery, Kafkas University Faculty of Medicine, Assoc. Dr., Kars, Turkey

²) Department of Cardiovascular Surgery, Kafkas University Faculty of Medicine, MD, Kars, Turkey

³) Department of Cardiovascular Surgery, Kafkas University Faculty of Medicine, Prof. Dr., Kars, Turkey

Abstract

Transthoracic gunshot injuries acquire immediate intervention with a correct approach as they have high mortality rates. The high mortality mainly depends on the presence of a probable concomitant injury of the cardiac, pulmonary and major thoracal vascular structures. In this case report, the emergent surgical management of a young female referred with a thoracal gunshot wound causing a massive pulmonary destruction is presented.

Keywords: Firearms; thorax; lung injury

*Başbuğ H. Z., Göçer H., Özişik K. Emergent operation of an extensive pulmonary damage after thoracal gunshot assault
EJCM 2017; 05 (4): 83-86. Doi: 10.15511/ejcm.17.00483.*

Introduction

Thoracic gunshot injuries are associated with high mortality due to concomitant pulmonary, cardiac or major vascular injuries.¹ The treatment of hemodynamically unstable patients with thoracic gunshot wounds is emergent due to major cardiac or vascular injuries. These patients should immediately proceed to the operating room.² In hemodynamically stable patients, diagnostic tests are needed before urgent transfer to the operation.³ In this paper, the emergent surgical management of a young woman with a transthoracic gunshot wound was presented. This case is diversified from others attributing to the presence of a pulmonary damage with a suspicious cardiac injury.

Case Report


A 22-year-old female referred with a thoracic gunshot wound. She was conscious with a blood pressure of 80/50 mmHg and heart rate of 124 bpm. She had fire-arm bullet entry wound just above her left nipple and an exit wound on her left posterior axillary line below the

Figure 1. Morphological view of the penetrating firearm wound. Note the contact muzzle stamp around the bullet entry hole.



sixth intercostal space. Physical examination of the entry wound revealed a contact muzzle stamp over the nipple (**Figure 1**). Posteriorly, there was a lateral bullet exit lesion between the sixth intercostal space and the fractured seventh rib. The Computerized Tomography (CT) Angiography of the thorax demonstrated the lack of great vessel injury and the presence of a massive left hemothorax consistent with the left pulmonary parenchymal injury (**Figure 1**). There was an interesting patchy opaque extravasation around the pericardium near the cardiac apex which might be related to a cardiac injury (**Figure 2**). Thus, the patient was taken to the operating room urgently. Following a median sternotomy, the pericardium was dissected 1 cm long to explore and see the pericardial

Figure 2. Computerized Tomography (CT) Angiography of the thorax demonstrates the presence of a massive left hemothorax with a contrast extravasation consistent with the left pulmonary parenchymal injury (A). The suspicious view in the CT Angiography implicating a cardiac injury (B).



aspirate. There was no hemopericardium consistent with a cardiac involvement that was suspected by the **Figure 1**. Left pleura was then widely opened beneath the sternum and 1200 ml of blood was aspirated from the left thorax. Upon examination of the left lung, the superior lobe was seriously injured. The parenchymal laceration was repaired with a 4-0 polypropylene by double running stitch (**Figure 1**). Near the bullet exit site over the inferior lobe, there was another parenchymal laceration which was also repaired with the same suturing technique. No postoperative complication happened.

Discussion

Advances in prehospital care systems enabled shorter delivery times for the penetrating trauma patients to the hospitals. “Scoop and run” strategy have replaced the older “stay and play” manner. This resulted in more trauma victims become more able to reach the emergency departments in extremis.⁴ Despite this advance in improvement in the transportation of the casualties to the hospital, overall survival after transthoracic gunshot injuries has not improved satisfactorily depending on the publications between 1966 and 2012.¹ From Bradley⁵ who reported the first case series in 1966 with a 20% of mortality, up to Okeye et al.¹ who declared an overall mortality rate of 78%, the survival after transthoracic gunshot wounds is still inefficiently low.

Transthoracic gunshot wounds are almost always associated with a hemodynamic compromise and require urgent appropriate intervention.⁶ However, hemodynam-

ically stable patients with transthoracic gunshot wounds would also have a potential risk for significant occult injury.⁷ Degiannis et al.⁸ reported 42% occult injuries among stable patients. Moreover, Richardson et al.⁹ observed occult injury in 63% of their patients who were initially stable. Thus, the traditional preoperative diagnostic tests in these instances become more crucial to determine the severity of the injury as well as to exclude a significant occult injury among the asymptomatic or less symptomatic cases. Intravenous contrast CT imaging is generally used for visualization the missile tract as well as to evaluate the vascular integrity.¹⁰ The use of CT-imaging as a screening tool is generally sufficient to rule out significant thoracic and mediastinal injury.¹¹

The operative approach in transthoracic penetrating trauma patients depends on the hemodynamic condition of the patient and the injury site.¹² In a hemodynamically stable patient with a mediastinal penetrating injury, the incision should be tailored meticulously. The majority of cardiac and thoracic injuries can comfortably be handled with a median sternotomy.¹³

However, for patients who arrived in extremis, lateral thoracotomy and aortic cross-clamping may be suitable to control the mediastinal bleeding rapidly.¹⁴ The surgical approach must be directed to the optimal exposure and easiest access. Injuries of the structures located in the posterior mediastinum including the thoracic esophagus, descending aorta, distal tracheobronchial structures need lateral thoracotomy while the cardiac and ascend-

Figure 3. The intraoperative image is demonstrating the pulmonary parenchymal destruction. Note the hemorrhage through the laceration site.



Figure 4. The intraoperative image is demonstrating the pulmonary parenchymal repair. Note the hemorrhage was stopped.



ing aorta injuries need median sternotomy.¹ In this case, a median sternotomy was preferred to evaluate the suspicious cardiac involvement that was seen on CT image (Figure 3). CT image revealed a suspicious contrast extravasation that may implicate a possible apical involvement as the bullet was glanced off. Upon performing a tiny pericardial incision, no pericardial blood was seen. By excluding the cardiac involvement, the left lung was

assessed and repaired through the sternotomy easily (Figure 4) and the drain tubes were successfully placed.

In conclusion, transthoracic firearm injuries are almost always associated with hemodynamic compromise and require rapid and appropriate resuscitative surgical exploration. Thus, depending on the injured structures and vital status of the patient, correct approach should be tailored in these life-threatening injuries.

References

1. Okoye OT, Talving P, Teixeira PG, Chervonski M, Smith JA, Inaba K, et al. Transmediastinal gunshot wounds in a mature trauma centre: Changing perspectives. *Injury* 2013;44(9):1198-203.
2. Stassen NA, Lukan JK, Spain DA, Miller FB, Carrillo EH, Richardson JD. Reevaluation of diagnostic procedures for transmediastinal gunshot wounds. *J Trauma* 2002;53(4):635-8.
3. Cornwell EE, Kennedy F, Ayad IA, Berne TV, Velmahos G, Asensio J, et al. Transmediastinal gunshot wounds: a reconsideration of the role of aortography. *Arch Surg* 1996;131(9):949-53.
4. Smith RM, Conn AK. Prehospital care-scoop and run or stay and play? *Injury* 2009;40(Suppl. 4):S23-6.
5. Bradley MN. Transmediastinal wounds. *Am Surg* 1966;32:847-52.
6. Kerimoğlu B, Köse S, Özışık K, Ertürk M, Koşar A, Orhan G. Yüksek hızlı ateşli silahlarla oluşan toraks yaralanmaları. *T Klin J Med Sci* 2001;21(4):249-52.
7. Nagy KK, Roberts RR, Smith RF, Joseph KT, An GC, Bokhari F, et al. Transmediastinal gunshot wounds: are "stable" patients really stable? *World J Surg* 2002;26:1247-50.
8. Degiannis E, Benn CA, Leandros E, Goosen J, Boffard K, Saadia R. Transmediastinal gunshot injuries. *Surgery* 2000;128:54-8.
9. Richardson JD, Flint LM, Snow NJ, Gray Jr LA, Trinkle JK. Management of transmediastinal gunshot wounds. *Surgery* 1981;90:671-6.
10. Grossman MD, May AK, Schwab CW, Reilly PM, McMahon DJ, Rondono M, et al. Determining anatomic injury with computed tomography in selected torso gunshot wounds. *J Trauma* 1998;45:446-56.
11. Hanpeter DE, Demetriades D, Asensio JA, Berne TV, Velmahos G, Murray J. Helical computed tomographic scan in the evaluation of mediastinal gunshot wounds. *J Trauma* 2000;49:689-94.
12. Peper WA, Obeid FN, Horst HM, Bivins BA. Penetrating injuries of the mediastinum. *Am Surg* 1986;52:359-65.
13. O'Connor JV, Scalea TM. Penetrating thoracic great vessel injury: impact of admission hemodynamics and preoperative imaging. *J Trauma* 2010;68:834-7.
14. Working Group, Ad Hoc Subcommittee on Outcomes, American College of Surgeons. Committee on trauma, practice management guidelines for emergency department thoracotomy. working group, ad hoc subcommittee on outcomes, American college of surgeons-committee on trauma. *J Am Coll Surg* 2001;193:303-9.

Received: 19/06/2017

Accepted: 21/11/2017

Published: 15/03/2017

Disclosure and conflicts of interest:

Conflicts of interest were not reported.

Corresponding author:

Hamit Serdar Başbuğ

Mail: s_basbug@hotmail.com

Which should be treated first; coronary artery disease or celiac artery compression syndrome?

İsmail Bıyık¹, Erol Bahtiyar², Nilgün Işıksaçan³

¹) Department of cardiology, Usak University, School of Medicine, Education and Research Hospital, Assoc. Dr., Usak, Turkey

²) Department of Cardiovascular Surgery, Usak University, School of Medicine, Education and Research Hospital, MD., Usak, Turkey

³) Department of Biochemistry, Bakirkoy Sadi Konuk Education and Research Hospital, MD., Istanbul, Turkey

Abstract

Celiac artery compression syndrome is rare. The differential diagnosis and the choice of therapeutic intervention in this syndrome are controversial. Herein, we report of a case of celiac artery compression syndrome and concomitant coronary artery disease and discuss differential diagnosis and treatment strategies.

Key words: Dunbar syndrome, celiac artery compression syndrome, coroner artery disease, intervention

Bıyık İ., Bahtiyar E., Işıksaçan N., Which should be treated first; coronary artery disease or celiac artery compression syndrome? EJCM 2017; 05 (4): 87-90. Doi: 10.15511/ejcm.17.00487.

Introduction

Celiac artery compression syndrome (CACS) is a rare clinical entity. It has also known as median arcuate ligament (MAL) or Dunbar syndrome. The disease results from the external compression of the celiac artery by the diaphragmatic crura or the median arcuate ligament.⁽¹⁾ The incidence of this rare condition has been reported from 1.76% to 4.0% in different radiologic studies.⁽¹⁾ Approximately, 85% of patients with CACS are asymptomatic.⁽¹⁾ The leading symptoms of this entity has been reported as abdominal pain (94%), postprandial abdominal pain (80%), nausea and vomiting (55.6%), weight loss (50%), bloating (39%), and abdominal pain triggered by exercise (8%).⁽²⁾ The differential diagnosis and the choice of therapeutic intervention in this syndrome are controversial.⁽²⁾ Herein, we report of a case of CACS and concomitant coronary artery disease (CAD) and discuss differential diagnosis and treatment strategies.

Case Report

Informed consent of presented patient has been taken for this report. A forty three years old male patient presented to our out-patient clinic of cardiology with the symptoms of long-standing epigastric pain, abdominal swelling and loss of appetite. His detailed history revealed that the symptoms increase especially after overdosed dinner, hiking and exercises. His habitual

history revealed excessive alcohol and cigarette consumption. His physical examination was unremarkable except abdominal obesity. His body mass index was 32 kg / m². His electrocardiogram and echocardiographic study were normal. His biochemical and laboratory investigations were in normal limits. Exercise electrocardiography showed horizontal ST segment depressions in leads DII, III and AVF at mid-level exercise.

Radionuclide myocardial perfusion scintigraphy also demonstrated reversible inferior wall hypo-perfusion and hypokinesia. Coronary angiography and selective celiac and mesenteric arteriography were performed for the differential diagnosis of symptoms. Coronary angiography displayed critical stenosis visually about 80 to 85% at the proximal level of well-developed obtuse margin branch of circumflex artery (**Figure 1**).

In celiac arteriography, there were an external compression on celiac artery causing critical stenosis visually about 80 to 85% and post-stenotic dilatation of the artery (**Figure 2**). Computed tomography angiography of abdominal arteries and reconstruction of data confirmed overhead external compression of celiac artery causing critical stenosis and post-stenotic dilatation of the artery (**Figure 3**). The case was discussed in the council of the cardiology and cardiovascular surgery of the institution. We decided to treat coronary artery disease as first intervention with the stenting of the ob-

Figure 1. Selective coronary angiography





Figure 2. Celiac angiography




tuse margin branch lesion of circumflex artery and to observe the patient at least one month for the symptoms relief. After successful stenting procedure (**Figure 4**), the symptoms of the patient quickly relieved, and he was discharged without symptoms. At first month visit, the patient was well condition and there were no his previous complaints.

Discussion

Symptomatic CACS is a very rare disease, which is clinically determined by the triad of postprandial abdominal pain, weight loss and an abdominal murmur because of the compression of the celiac artery.⁽³⁾ Fibrous median arcuate ligament usually bestrides the celiac artery at the level of the first lumbar vertebra in 10% to 24% of the general population but this entity causes symptoms a few of these patients.⁽⁴⁾

There are a lot of reports on this syndrome in literature but its association with coronary artery disease has not been established to date. In the presented case, we took in consideration both CAD and CACS for the source and cause of complaints. Investigative work up was performed in two directions. Celiac arteriography added to coronary angiographic study revealed both critical coronary artery stenosis and celiac artery compression.

Figure 3. Computed tomography angiography of abdominal arteries and reconstruction of data



The case was discussed in details in institutional council of cardiology and cardiovascular surgery. Because CACS is frequently encountered but rarely causes symptoms, we decided to treat CAD at first. After successfully performed coronary intervention, patient's complaints and symptoms were disappeared.

CACS could be investigated with Doppler ultrasonography, computed tomography angiography, conventional angiography and magnetic resonance angiography. The traditional method for the treatment of CACS syndrome is open surgical intervention. Recent reports suggest that the laparoscopic surgery is best approach in the treatment of CACS.⁽⁵⁾ Laparoscopic MAL release has some advantageous such as decreased postoperative pain, lower incidence of postoperative adhesions, shorter recovery time and hospital stay period.⁽⁵⁾ Nowadays, this method has become standard surgical approach for the treatment of CACS.⁽²⁾

In addition, complex reconstruction procedures such as patch angioplasty, aorta-celiac bypass, reanastomosis of celiac artery to aorta have been performed when needed.⁽²⁾ In patients having persistent symptoms and residual critical celiac artery stenosis after laparoscopic MAL release, percutaneous endo-

Figure 4. Stenting of obtuse marginal branch of circumflex artery



vascular intervention and stenting of persistent critical celiac artery stenosis may be performed.⁽⁶⁾ CAD may mimic the symptoms of CACS especially such as associated with the exercise. Investigational work up should include both diseases.

Conclusion

Because symptomatic CACS is very rare entity, CAD should be taken into account. Diagnostic work up should include CAD. It may be a reasonable approach to treat coronary artery disease primarily in terms of clinical and prognostic aspects.

References

1. Gümüş H, Gümüş M, Tekbaş G, Önder H, Ekici F, Çetinçakmak MG, Bilici A. Clinical and multidetector computed tomography findings of patients with median arcuate ligament syndrome. *Clin Imaging*. 2012; 36(5): 522-5.
2. Kim EN, Lamb K, Relles D, Moudgill N, DiMuzio PJ, Eisenberg JA. Median Arcuate Ligament Syndrome-Review of This Rare Disease. *JAMA Surg*. 2016; 151(5): 471-7.
3. Kuruvilla A, Murtaza G, Cheema A, Arshad HMS. Median Arcuate Ligament Syndrome: It Is Not Always Gastritis. *J Investig Med High Impact Case Rep*. 2017; 5(3): 2324709617728750.
4. Duffy AJ, Panait L, Eisenberg D, Bell RL, Roberts KE, Sumpio B. Management of median arcuate ligament syndrome: a new paradigm. *Ann Vasc Surg*. 2009; 23: 778-784.
5. Duran M, Simon F, Ertas N, Schelzig H, Floros N. Open vascular treatment of median arcuate ligament syndrome. *BMC Surg*. 2017; 17(1): 95.
6. Balkan A, Alkan S, Yidirim AE, Konduk BT, Er RE, Gulsen MT. A Rare Cause of Chronic Abdominal Pain: Median Arcuate Ligament Syndrome. *Bozok Med J* 2017; 7(3): 106-9.

Received: 16/05/2017

Accepted: 21/10/2017

Published: 15/12/2017

Disclosure and conflicts of interest:

Conflicts of interest were not reported.

Corresponding author:

Dr. İsmail Biyık

Mail: ismailbiyikmd@yahoo.com

It's not fair to compare transcatheter aortic valve implantation with surgical replacement

Taylan Adademir¹, Davut Çekmecelioğlu², Deniz Çevirme¹, Mehmet Kalender³, Kamil Boyacıoğlu⁴, Fuat Büyükbayrak⁵, Hızır Mete Alp⁶

¹ Kartal Kosuyolu Yüksek İhtisas Research and Training Hospital, Cardiovascular Surgery Clinic, M.D., Istanbul, Turkey

² Kartal Kosuyolu Yüksek İhtisas Research and Training Hospital, Cardiovascular Surgery Clinic, Assis. Dr., Istanbul, Turkey

³ Kocaeli University of Health Sciences, Derince Education and Research Hospital, Cardiovascular Surgery Clinic, Op. Dr., Kocaeli, Turkey

⁴ Istanbul Bağcılar Research and Training Hospital, Cardiovascular Surgery Clinic, Op. Dr., Istanbul, Turkey

⁵ Izmit Konak Hospital, Cardiovascular Surgery Clinic, Assoc. Dr., Izmit, Turkey

⁶ Okan University Hospital, Cardiovascular Surgery Clinic, Prof. Dr., Istanbul, Turkey

Abstract

Surgical aortic valve replacement (SAVR) is still class I recommendation for the treatment of patients with symptomatic severe aortic valve stenosis (AS) according to recent guidelines. Although it assumed as one of the most successful intervention of cardiac surgery; currently the transcatheter technique (transcatheter aortic valve replacement-TAVR) has come into notice as a feasible and effective option to treat high risk or inoperable aortic stenosis patients. New reports have been publishing comparison of these two techniques by numbers increasing with this technology¹. On the stage, we would like to discuss about fairness of the comparison mentality in terms of a feature-by-feature style.

Key words: Dunbar syndrome, celiac artery compression syndrome, coroner artery disease, intervention

Adademir T., Çekmecelioğlu D., Çevirme D., Kalender M., Boyacıoğlu K., Büyükbayrak F., Alp H. M. It's not fair to compare transcatheter aortic valve implantation with surgical replacement EJCM 2017; 05 (4): 91-94. Doi: 10.15511/ejcm.17.00491.

As we all know, gold standard therapy should be surgical replacement for the patients with severe calcific aortic stenosis echocardiographically, either symptomatic or asymptomatic. Conventional SAVR yields excellent results after 1 year in lower-risk patients. Despite that, TAVR is also recommended as class I indication on recent guidelines for patients who inoperable or have prohibitive high surgical risk with last decision given to Heart Valve Team.²

Nonetheless, constitutional complications of TAVR have been surfacing and limiting its use such as embolization of calcified debris and consequent cerebral infarction; complete AV block and need of pacemaker insertion; paravalvular leakage and its effect on long-term survival; coronary ostium closure and even aortic rupture.^{3,4} Up to 42.5% complete AV block, 15% moderate-to-severe paravalvular leak, and 84% new cerebral infarction have been reported after TAVR.⁵

Despite higher complication rates in terms of major adverse cardiac and cerebrovascular events (MACCE) among TAVR patients; early studies those compared SAVR and TAVR revealed that TAVR is not inferior to surgical AVR in mortality rates. And even more recent trials claims superior mortality results with TAVR. Randomised and observational evidence adjusted on the baseline patients' characteristics finds a similar risk for 30 days mortality, 1-year mortality, stroke, myocardial infarct (MI) and acute renal injury in TAVR and SAVR.⁶ Also Vernat et al⁷ reported that mitral annular geometry is preserved better by TAVR than by SAVR. They concluded that TAVR might be more physiologic approach for aortic replacement. Kocaaslan et al⁸ found that the increase in terms of quality of life parameters in the TAVR group was greater than the AVR group at the end of the 3rd month. Probably, the best attribution to TAVR is being non-invasive method that no need for cardiopulmonary bypass (CPB) during the procedure.

On the other hand, patients with high surgical risk did not have any significant difference for the first year mortality according to two large registries (Partner and GARY = German Registry). But reports showed a two-fold increased risk of cerebrovascular accident in the TAVR group. The PARTNER A and B trials showed that

survival has been remarkably good, but stroke and perivalvular leakage require further device development.⁹ These findings are far from being gold standard. However, progressing technology will continue to seek new answers to these problems. So, can not surgeons replace the aortic valve surgically which is gold standard with low mortality rates, even for high risk patients?

Fairness always needs feature-by-feature comparison. High technology should be compared with high technology. By using higher technology valves, for example suturless valves, we can speed up the surgery. By reducing cross clamp times, we would decrease operative mortality and morbidity. By using high technology, we can perform aortic valve surgery through minimal approaches via mini (CBP) circuits which would decrease morbidity of the pathology. Mini CBP circuits and cannulas may also decrease mortality rates by reducing inflammatory response which may yield sepsis condition.

Sutureless AVR exploitation self-expanding bioprosthesis is a new and favourable alternative to conventional surgical AVR in elderly and high-risk surgical patients. The advanced benefits of this technology involve improved implantability, shorter aortic cross-clamp and cardiopulmonary bypass (CPB) times, promised hemodynamic presentation and easier approach for minimally invasive surgery.¹⁰

Does faster replacement of the valve make a real significant effect especially in recent days which myocardial protection methods are well developed? There are several publications seeking answers to this question. Ranucci et al¹¹ showed that the aortic cross clamping time was an independent predictor of severe cardiovascular morbidity, with an increased risk of 1.4% per 1 min increase. Patients with a left ventricular ejection fraction $\leq 40\%$ showed the most relevant clinical benefits induced by reduction in cross clamp time.

Additionally, this technology admits entire removal of the diseased native valve and also consists a appropriate alternative to multiple valve procedures or associated coronary artery bypass grafting. Several European case series have demonstrate excellent early clinical and hemodynamic results.¹²

We can observe zero in hospital mortality and significantly lower paravalvular leak in high risk patients whom hospital mortality is 5.3% for TAVR group by adding only one high technology to surgical arm which is sutureless valve. Comparative reports between sutureless valves and TAVR in intermediate and high risk patients have shown lower rate of perioperative complications in terms of MACCE and enhanced survival at 24-month follow-up with sutureless valves.^{3,13} We can easily assume that the results would be even better if we can add mini CBP techniques into these technology. Consequently, sutureless aortic bioprostheses has been situated as an alternative to conventional surgical AVR or TAVR in elderly and high-risk patients. However, the need of multi-centered, randomized trials can not be ignored.

Cardiovascular surgeons involved in the treatment of valve diseases should be open to the introduction of innovative methods, technologies or hypothesis. Also he/she should not get carried off and should always examine the outcomes in the light of well-conducted short and long term clinical trials. Apparently that we must utilize the science strictly to keep patients' safety.¹⁴

TAVR is growing as a minimally invasive therapy

for patients with severe AS. Today, it is completely acceptable that TAVR is already preferable in an increasing proportion of elderly and high risk patients; nevertheless, this procedure has potential for serious complications. The recent evidence and outcomes has showed that its feasibility and efficacy. They draw the attention of industry and physicians to this new technology. There is little evidence on the long-term outcomes in spite of short-term efficacy is good. We aware that AVR treats the disease quite constantly and it is still considered the standard treatment for most of patients. The future is expected to be more wealthy as new developments and statistics got from continued trials. They will also provide the background to expand physicians' decision area and their applications. Lastly, Bavaria¹⁵ et al found that not only were surgeons actively involved in the treatment decision-making process, but also played a significant role in the valve procedure, including deployment and post-procedural care of TAVR patients according the Society of Thoracic Surgeons(STS) survey. That is why cardiac surgeons need to have real and fair statement about TAVR.

Choice of SAVR, TAVR or sutureless AVR and indications would be concisely established by fairness with a broader spectrum of patients.

References

1. Leon, Martin B., et al. "Transcatheter or surgical aortic-valve replacement in intermediate-risk patients." *N Engl J Med* 2016.374 (2016): 1609-1620.
2. Nishimura, Rick A., et al. "2014 AHA/ACC guideline for the management of patients with valvular heart disease." *Circulation* (2014):
3. Biancari, Fausto, et al. "Immediate outcome after sutureless versus transcatheter aortic valve replacement." *Heart and vessels* 31.3 (2016): 427-433.
4. Daneault, Benoit, et al. "Stroke associated with surgical and transcatheter treatment of aortic stenosis: a comprehensive review." *Journal of the American College of Cardiology* 58.21 (2011): 2143-2150.
5. Sinning, Jan-Malte, et al. "Aortic regurgitation index defines severity of peri-prosthetic regurgitation and predicts outcome in patients after transcatheter aortic valve implantation." *Journal of the American College of Cardiology* 59.13 (2012): 1134-1141.
6. Nagaraja, Vinayak, et al. "Transcatheter versus surgical aortic valve replacement: a systematic review and meta-analysis of randomised and non-randomised trials." (2014): e000013.)
7. Vergnat, Mathieu, et al. "The effect of surgical and transcatheter aortic valve replacement on mitral annular anatomy." *The Annals of thoracic surgery* 95.2 (2013): 614-619.
8. Kocaaslan, Cemal, et al. "Comparison of Transcatheter Aortic Valve Implantation versus Surgical Aortic Valve Replacement to Improve Quality of Life in Patients > 70 Years of Age with Severe Aortic Stenosis." *Brazilian journal of cardiovascular surgery* 31.1 (2016): 1-6.
9. Svensson, Lars G., et al. "A comprehensive review of the PARTNER trial." (2013): S11-S16.
10. Santarpino G, Pfeiffer S, Concistré G, Grossmann I, Hinzmann M, Fischlein T. The Perceval S aortic valve has the potential of shortening surgical time: does it also result in improved outcome? *Ann Thorac Surg.* 2013;96(1):77-81.
11. Ranucci, Marco, et al. "Aortic cross-clamp time, new prostheses, and outcome in aortic valve replacement." *The Journal of heart valve disease* 21.6 (2012): 732-739.
12. Folliguet TA, Laborde F, Zannis K, Ghorayeb G, Haverich A, Shrestha M. Sutureless Perceval aortic valve replacement: results of two European centers. *Ann Thorac Surg.* 2012;93(5):1483-8
13. Santarpino G, Pfeiffer S, Jessl J, Dell'Aquila AM, Pollari F, Pauschinger M, et al. Sutureless replacement versus transcatheter valve implantation in aortic valve stenosis: a propensity-matched analysis of 2 strategies in high-risk patients. *J Thorac Cardiovasc Surg.* 2014;147(2):561-7
14. Saadi, Eduardo Keller. "Percutaneous aortic aortic valve replacement: myth or reality?." *Brazilian Journal of Cardiovascular Surgery* 23.1 (2008): 93-96.
15. Bavaria, J. E., Prager, R. L., Naunheim, K. S., Allen, M. S., Higgins, R. S., Thourani. STS Executive Committee. (2017). Surgeon Involvement in Transcatheter Aortic Valve Replacement in the United States: A 2016 Society of Thoracic Surgeons Survey. *The Annals of Thoracic Surgery.*

Received: 30/07/2017

Accepted: 09/11/2017

Published: 15/12/2017

Disclosure and conflicts of interest:

The authors declare no conflict of interest.

Corresponding author:

Davut Çekmecelioglu

Mail: d.cekmecelioglu@yahoo.com

E-Journal of Cardiovascular Medicine welcomes scientific contributions in the field of cardiovascular and thoracic surgery - all aspects of surgery of the heart, vessels and the chest in various article types: new ideas, brief communications, work in progress, follow-up studies, original articles, best evidence topics, case reports, reports on unexpected results etc. All manuscripts shall be reviewed by the Editor-in-Chief, Associate Editors, Invited Referees and a Statistician when appropriate. If accepted, articles will be posted online and opened up for discussion. Acceptance criteria are based on the originality, significance, and validity of the material presented.

All material to be considered for publication in E-Journal of Cardiovascular Medicine should be submitted in electronic form via the journal's online submission system. (<http://my.ejmanager.com/ejcm/>)

A cover letter should be enclosed to all new manuscripts (to be filled in online), specifying the name of the journal and the type of paper, and including the following statements:

- The manuscript should not be previously published in print or electronic form and is not under consideration by another publication.
- All authors should contribute to the content of the article.
- All authors should read and approve the submission of the manuscript to ICVTS.
- Subject to acceptance, authors will sign an exclusive license to publish.
- No ethical problem or conflict of interest should exist.

If your first language is not English, we recommend that you consult an English language editing service to ensure that the academic content of your paper is fully understood by journal editors and reviewers. Language editing does not guarantee that your manuscript will be accepted for publication.

Manuscripts should be prepared using a word-processing package (save in .doc, .docx or .rtf format). The font type and font size should preferably be Arial or Times New Roman 11 points. The manuscript should be double-spaced and should include line and page numbers. The lines of the reference list do not need to be numbered; include a section break before.

Manuscripts should be organized as follows:

(a) Title page; (b) Abstract and Key words; (c) Text with the following sections (not applicable for article types with unstructured abstracts): Introduction, Materials and Methods, Results, Discussion, Acknowledgement (optional), Funding statement, Conflict of interest statement, (d) Figure (and Video) legends; (e) Tables; (f) References.

Title page (1st page): Title: should be brief and descriptive (100 characters) - no abbreviations are allowed, even if well known.

Authors: list all authors by full first name, initial of or full middle name and family name. Qualifications are not required. Ensure the author names correspond (in spelling and order of appearance) with the metadata of the system

Institution(s): include the name of all institutions with the location

(department, institution, city, country) to which the work should be attributed (in English). Use superscript numbers to connect authors and their department or institution.

Corresponding author: The full name, full postal address, telephone/fax numbers and the e-mail address should be typed at the bottom of the title page.

Meeting presentation: If the manuscript was (or will be) presented at a meeting, include the meeting name, venue, and the date on which it was (or will be) read; also indicate if you have submitted an Abstract of this manuscript for the EACTS or ESTS annual meeting and whether it has been accepted (if known).

Word count: The total number of words of the whole article (including title page, abstract, main text, legends, tables and references) must be specified on the title page.

Abstract (2nd page): An abstract should be a concise summary of the manuscript. Reference citations are not allowed. The abstract should be factual and free of abbreviations, except for SI units of measurement.

Keywords: Following the abstract, 3-6 keywords should be given for subject indexing.

Introduction: Should state the purpose of the investigation and give a short review of pertinent literature.

Materials and methods: Should be described in detail with appropriate information about patients or experimental animals. Use of abbreviations renders the text difficult to read; abbreviations should be limited to SI units of measurement and to those most commonly used, e.g. VSD, ASD, CABG (abbreviations should not be included in headings and extensions should be included at first mention).

Results: Results should be reported concisely and regarded as an important part of the manuscript. They should be presented either in tables and figures, and briefly commented on in the text, or in the text alone. Repetition of results should be avoided!

Discussion: The discussion is an interpretation of the results and their significance with reference to pertinent work by other authors. It should be clear and concise.

Acknowledgement: Acknowledgements and details of non-financial support must be included at the end of the text before the references. Personal acknowledgements should precede those of institutions or agencies.

Tables: All tables must be included in the manuscript file, should start on separate pages and be accompanied by a title, and footnotes where necessary. The tables should be numbered consecutively using Arabic numerals. Units in which results are expressed should be given in parentheses at the top of each column and not repeated in each line of the table.

References: Authors are responsible for checking the accuracy of all references. If you use EndNote or Reference Manager to facilitate referencing citations (not required for submission), this journal's style is available for use. References should be numbered in order of appearance in the text (in Arabic numerals in parentheses) and must be listed numerically in the reference list. Journal titles and author initials should be properly abbreviated and punctuated.

GENERAL RULES

Files should be prepared as a Word document using font size 12 Times New Roman characters, double-spaced and with 2.5 cm margins on each side, top and bottom. Only standard abbreviations should be used; other shortened phrases should be indicated in parentheses as used in the text. Generic or chemical names of drugs should be used instead of trade names.

ETHICAL ISSUES

Publishing responsibilities of authors and Ethics

The publication of an article in a peer-reviewed journal is an essential building block in the development of a coherent and respected network of knowledge. It is a direct reflection of the quality of work of the author and the institutions that support them. Peer-reviewed articles support and embody the scientific method. It is therefore important to agree upon standards of expected ethical behavior.

Reporting standards

Authors of reports of original research should present an accurate account of the work performed as well as an objective discussion of its significance. Underlying data should be represented accurately in the paper. A paper should contain sufficient detail and references to permit others to replicate the work. Fraudulent or knowingly inaccurate statements constitute unethical behavior and are unacceptable. Review and professional publication articles should also be accurate and objective, and editorial 'opinion' works should be identified as such.

Hazards and human or animal subjects

If the work involves chemicals, procedures or equipment that have any unusual hazards inherent in their use, the author must clearly identify these in the manuscript. If the work involves the use of animal or human subjects, the author should ensure that the manuscript contains a statement that all procedures were performed in compliance with relevant laws and institutional guidelines and that the appropriate institutional committee(s) has approved them. Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

Use of patient images or case details

Studies on patients or volunteers require ethics committee approval and informed consent, which should be documented in the paper. Appropriate consents, permissions and releases must be obtained where an author wishes to include case details or other personal information or images of patients and any other individuals in publication. Written consents must be retained by the author and copies of the consents or evidence that such consents have been obtained must be provided to us on request. Particular care should be taken with obtaining consent where children are concerned (in particular where a child has special needs or learning disabilities), where an individual's head or face appears, or where reference is made to an individual's name or other personal details.

Originality and plagiarism

The authors should ensure that they have written entirely original works, and if the authors have used the work and/or words of

others, that this has been appropriately cited or quoted. Plagiarism takes many forms, from 'passing off' another's paper as the author's own paper, to copying or paraphrasing substantial parts of another's paper (without attribution), to claiming results from research conducted by others. Plagiarism in all its forms constitutes unethical publishing behavior and is unacceptable.

Data access and retention

Authors may be asked to provide the raw data in connection with a paper for editorial review, and should be prepared to provide public access to such data (consistent with the ALPSP-STM Statement on Data and Databases), if practicable, and should in any event be prepared to retain such data for a reasonable time after publication.

Multiple, redundant or concurrent publication

An author should not in general publish manuscripts describing essentially the same research in more than one journal or primary publication. Submitting the same manuscript to more than one journal concurrently constitutes unethical publishing behavior and is unacceptable. In general, an author should not submit for consideration in another journal a previously published paper. Publication of some kinds of articles (e.g. clinical guidelines, translations) in more than one journal is sometimes justifiable, provided certain conditions are met. The authors and editors of the journals concerned must agree to the secondary publication, which must reflect the same data and interpretation of the primary document. The primary reference must be cited in the secondary publication.

Acknowledgement of sources

Proper acknowledgment of the work of others must always be given. Authors should cite publications that have been influential in determining the nature of the reported work. Information obtained privately, as in conversation, correspondence, or discussion with third parties, must not be used or reported without explicit, written permission from the source. Information obtained in the course of confidential services, such as refereeing manuscripts or grant applications, must not be used without the explicit written permission of the author of the work involved in those services.

Fundamental errors in published works

When an author discovers a significant error or inaccuracy in his/her own published work, it is the author's obligation to promptly notify the journal editor or publisher and cooperate with the editor to retract or correct the paper. If the editor or the publisher learns from a third party that a published work contains a significant error, it is the obligation of the author to promptly retract or correct the paper or provide evidence to the editor of the correctness of the original paper.

Authorship of the paper

Authorship should be limited to those who have made a significant contribution to the conception, design, execution, or interpretation of the reported study. All those who have made significant

contributions should be listed as co-authors. Where there are others who have participated in certain substantive aspects of the research project, they should be acknowledged or listed as contributors. The corresponding author should ensure that all appropriate co-authors and no inappropriate co-authors are included on the paper, and that all co-authors have seen and approved the final version of the paper and have agreed to its submission for publication.

Changes to authorship

This policy concerns the addition, deletion, or rearrangement of author names in the authorship of accepted manuscripts. Before the accepted manuscript is published in an online issue:

Requests to add or remove an author, or to rearrange the author names, must be sent to the Journal Manager by the corresponding author of the accepted manuscript, and must include:

The reason the name should be added or removed, or the author names rearranged

Written confirmation (e-mail, fax, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed

Requests that are not sent by the corresponding author will be forwarded by the Journal Manager to the corresponding author, who must follow the procedure described above.

Note that:

- Journal Managers will inform the Journal Editors of any such requests
- Publication of the accepted manuscript in an online issue is suspended until authorship has been agreed
- After the accepted manuscript has been published in an online issue:

Any requests to add, delete or rearrange author names in an article published in an online issue will follow the same policies as noted above and may result in a corrigendum.

TYPES OF PAPERS

Original Articles

Original articles should consist of sections titled as “Abstract, Introduction, Materials and Methods, Results, Discussion and Conclusion”. For information about the abstract, refer to ‘Manuscript Formatting’ section.

The Introduction section of the manuscript should clearly state the purpose of the manuscript and include a brief summary of the most relevant national and international literature stating the main purposes and research question of the study. Contradictory aspects of the research, if present, should be mentioned. The expected contribution of this study to family medicine and practice should be highlighted.

The Materials and Methods section should describe the study population and the study design, with adequate information on the

techniques, materials and methods used. The section should include information of the study type, population, sample, sample size and selection of the sample. Validity and reliability of scales and questionnaires used also should be referred to. A clear description of the statistical methods should also be given.

The Results section should include a detailed report on the findings of the study. All figures, tables and illustrations should be used in this section. Results should be presented either as text or figures and/or tables and not be replicated.

The Discussion section of the study should emphasize the importance of the results and compare them with the results of other authors with relevant citations from the most recent literature. Study limitations and strengths should be specified. Suggestions for further studies in this area should be added.

The Conclusion should include the main conclusions based on the results of the research, emphasize the contributions of the study to family practice and propose original suggestions. A brief revision of all the results and the discussion should be avoided.

Original articles excluding case reports and systematic reviews should not exceed 3000 words excluding the abstract, references and tables. Case reports should not exceed 1000 words excluding the abstract, references and tables. There are no restrictions for systematic reviews.

Short Reports

Short Reports are accepted when the research topic, aim and results of the study are in limited in scope and in cases that do not require writing a full original article. Short Reports can be described as a summarized version that have been prepared according to the structure of research articles. Publishing an article as a short report does not reflect a lower quality. The same rules as relevant to original articles apply to preparing a short report, but structured abstracts are not mandatory references and tables should not exceed 6 and 2 in number, respectively. Abstracts should not exceed 100 words and the text should be restricted to a maximum of 1000 words.

Reviews

Reviews are evidence-based articles about a specific topic using relevant citations from the most recent literature with the authors’ conclusions on this subject. The author is expected to have conducted research on the subject and to have experience in order to discuss and analyze the subject. There is no obligation to follow a particular format and may contain subtitles depending on the subject. The text should not exceed 4000 words excluding the title, abstracts, references and tables. E Journal of Cardiovascular Medicine, only publishes review articles solicited by the editors.

Letters to Editor and Comments

Letters to the editor or comments can be sent to provide commentary and analysis concerning an article published in the journal, to give information about ongoing research, to provide informa-

tion in cardiology and cardiovascular-vascular-endovascular surgery, cardio-metabolic and vascular sciences. Letters to the editor or comments may include an optional title, tables and references. These articles should not exceed 1000 words.

What Would You Do?

These are brief articles discussing cases and situations encountered in cardiology and cardiovascular surgery with a biopsychosocial approach. If necessary, photographs (with permission from the patient/owner) may be added. Sections should consist of a title, case report, discussion, questions and answers. Brief comments can be sent to provide commentary on previous articles and case reports written by other authors. Comments should include the number of the journal the article was published in. The text should not exceed 1000 words.

International Reprints

Translations of important documents, declarations and guidelines prepared by international organizations in the field of cardiology and cardiovascular surgery, may be published in the journal. Presubmission Inquiry to the Editorial Board of the Journal before submitting the article is recommended. It is the translator's responsibility to obtain permission from the owner of the original manuscript for publication and translation.

News

These articles focus on advances and innovations in clinical topics relevant to cardiology and cardiovascular surgery. There is no obligation to follow a particular format. The text should be limited to 1000 words.

Editorials

Editorials usually provide information about the editorial policy of E Journal of Cardiovascular Medicine, give commentary and feedback on articles published in the journal, draw attention to topics of current interest and give information related to and discuss the development of cardiology and cardiovascular surgery in the world. They are mainly written by the members of the Editorial Board. Editorials are limited to 2000 words with some exceptions and may include a title and references when necessary.

MANUSCRIPT FORMATTING

Manuscripts should be designed in the following order:

Title page

Abstract

Main text

References

Tables, figures and illustrations

Title Page

The title page of the manuscript should include: The title, first

and last names of each author. Complete affiliation and title for each author, with the name of department (s) and institution (s) to which the work should be attributed.

The corresponding author should be clearly identified with name, address, telephone- facsimile number and email address for correspondence about the manuscript. Authors should clearly indicate if the article has previously been presented at a congress or scientific meeting. The title should be concise and informative without abbreviations and not exceed 10 words.

Abstract

Abstracts should be exact in English, with a minimum of 150 and maximum of 350 words. Abstracts of original research articles should be structured under subheadings as follows: objectives, methods, results and conclusion. A maximum of 3 key words should be added to English abstracts.

Text

The text contains the rest of the manuscript. It is structured differently according to the type of manuscript (original research article, review, etc.). For example, original research articles should consist of aim and objectives, methods, results, discussion and conclusion.

References

References should be cited in consecutive numerical order as first mentioned in the text and designated by the reference number in parentheses. If the number of authors for the reference is more than 6 authors, list the first three authors and add "et al".

Journal names should be abbreviated as used in Index Medicus. References should be cited in the Vancouver style. For detailed information please visit the relevant link

Examples:

For research articles follow the example below:

– Verschuren WM, Jacobs DR, Bloemberg BP, et al. Serum total cholesterol and long-term coronary heart disease mortality. JAMA 1995; 274(2): 131–6.

For book chapters follow the example below:

– Rakel RE. The family physician. In: Rakel RE, editor. Textbook of family practice. 5th ed. Philadelphia: W.B. Saunders; 1995. p. 3-19.

For web pages follow the example below:

– Guidance for clinicians. An International Benchmarking Study. <http://www.who.int/topics/surgery/> accessed: 29/09/2002.

Tables and Figures

Legends should take place on the top of the page for tables, and bottom of the page for figures and placed on separate pages. Explain all nonstandard abbreviations in footnotes.



E Journal of Cardiovascular Medicine
is a global e-journal targeting articles on:

- clinical cardiology,
- interventional cardiology,
 - arrhythmia,
- cardiovascular surgery,
- vascular & endovascular surgery,
- vascular biology